

Associative-algebraic approach to logarithmic conformal field theories

N. Read ^a and Hubert Saleur ^{b,c}

^a *Department of Physics*

Yale University, P.O. Box 208120, New Haven, CT 06520-8120 USA

^b *Service de Physique Théorique, CEA Saclay,
Gif Sur Yvette, 91191, France*

^c *Department of Physics and Astronomy, University of Southern California
Los Angeles, CA 90089, USA*

Abstract

We set up a strategy for studying large families of logarithmic conformal field theories by using the enlarged symmetries and non-semi-simple associative algebras appearing in their lattice regularizations (as discussed in a companion paper). Here we work out in detail two examples of theories derived as the continuum limit of XXZ spin-1/2 chains, which are related to spin chains with supersymmetry algebras $gl(n|n)$ and $gl(n+1|n)$, respectively, with open (or free) boundary conditions in all cases. These theories can also be viewed as vertex models, or as loop models. Their continuum limits are boundary conformal field theories (CFTs) with central charge $c = -2$ and $c = 0$ respectively, and in the loop interpretation they describe dense polymers and the boundaries of critical percolation clusters, respectively. We also discuss the case of dilute (critical) polymers as another boundary CFT with $c = 0$. Within the supersymmetric formulations, these boundary CFTs describe the fixed points of certain nonlinear sigma models that have a supercoset space as the target manifold, and of Landau-Ginzburg field theories. The submodule structures of indecomposable representations of the Virasoro algebra appearing in the boundary CFT, representing local fields, are derived from the lattice. A central result is the derivation of the fusion rules for these fields.

1 Introduction

The interest in logarithmic conformal field theories (CFTs) (see [1] for an introduction) has grown over the last ten years or so as their potential role in condensed matter as well as string theory applications has become more evident. Condensed matter applications include critical geometrical models in two dimensions such as percolation and polymers (see [2] for a recent contribution and review), and critical points in non-interacting disordered fermion models in 2+1 dimensions (see [3, 4, 5] for recent contributions), such as the transition between plateaux in the integer quantum Hall and spin quantum Hall effect. String theory applications include the study of sigma models on non-compact (super-) target spaces such as $PSU(1,1|2)$ [6]. Logarithmic CFTs are also fascinating mathematical objects in their own right, bound to attract the attention of representation theory experts.

Logarithmic CFTs were encountered as early as 1987 by Knizhnik [7]. In 1992, Rozansky and Saleur tackled in some details one of the simplest such theory, the WZW model on the $U(1|1)$ supergroup. They found that the non-semisimplicity of $U(1|1)$ carried over to the CFT, with the possibility of a non-diagonal Virasoro generator (here, proportional to the superalgebra Casimir) L_0 , indecomposable operator product expansions (OPEs), and logarithms in some of the correlators [8]. The potential generality of these features was realized almost simultaneously by Gurarie in 1993 [9], who coined the name logarithmic CFT.

Certainly the theory of symplectic fermions is the best understood logarithmic CFT [10]. [A close second is the WZW model on $U(1|1)$ where powerful results were obtained recently using the

minisuperspace approach [11]. It is not clear to what extent the current algebra structure makes this example qualitatively different from the logarithmic CFTs of more physical interest.] A natural extension of the symplectic fermions has given rise to the so-called “augmented $c_{p,1}$ models”, which exhibit an enlarged triplet $\mathcal{W}$ algebra [10, 12, 13]. An essential feature of these theories is that the identity has a logarithmic partner, and thus that there are at least two fields of vanishing dimension. There have been some attempts to generalize the structure of the augmented $c_{p,1}$ models to other series, in particular the $c_{p,2}$ case [14, 15].

One of the most interesting cases, with numerous condensed matter applications to disordered systems, corresponds to vanishing central charge. Consideration of singularities in OPEs as $c \rightarrow 0$ led Gurarie to the conjecture that in these theories the stress energy tensor T had a logarithmic partner t such that L_0 had a Jordan cell of dimension 2 at the level $h = 2$ [16]. In this and in subsequent papers Gurarie and Ludwig [17, 18] proposed to use the modes of t to define an extension of the Virasoro algebra characterized by a parameter b . [Although appealing intuitively and successful in some cases, the idea has not borne much fruit, in particular because of considerable difficulties in putting together chiral and non chiral sectors.] The problem has also been discussed from a slightly different point of view of the replica limit by Cardy [19].

Most of the approaches mentioned so far start from the consideration of abstract conformal field theories. Such generality may not be the easiest way to proceed. In 2001, we defined families of supersymmetric lattice models, whose continuum limits we identified with the IR limits of conformal supersymmetric non-linear sigma models, or in some cases with Landau-Ginzburg theories [20]. [For instance, one of these lattice models was defined using alternating fundamental and conjugate representations of $\mathfrak{sl}(n+1|n)$ and argued, in the continuum limit, to be described by the IR limit of the $CP^{n|n}$ model at $\theta = \pi$. This model is closely related to the properties of hulls of percolation [3]. A similar model was defined for dilute and dense polymers.] In all these cases, we obtained the full spectrum of the associated logarithmic conformal field theory, and found considerable richness, including an infinite set of Virasoro primary fields which all have rational conformal weights, and indications of a large non chiral symmetry algebra commuting with Virasoro. We note that in all our examples with $c = 0$, there was a single field with vanishing dimension. We expect in general that the logarithmic theories we uncovered do not coincide with any of the extended $c_{p,1}$ or $c_{p,2}$ models mentioned before (except for the symplectic fermions).

It is worthwhile to consider a broad view of the strategies pursued in these various references. First, we point out that the traditional idea of a *rational* CFT contains two essential ingredients [21]: the local fields (or corresponding states) fall into a *finite* number of representations of the chiral algebra (which in the strictest view of rational CFT is generated by a finite number of bosonic local fields of integer conformal weights), and these representations are semisimple, that is fully decomposable as direct sums of irreducible representations of the chiral algebra. Hence also the operator products of local fields decompose fully into sums of these local fields. Next, we note that the occurrence of logarithms in correlators is due to L_0 not being diagonalizable, and this results from the appearance of fields transforming in non-semisimple representations of the Virasoro algebra. One is then led to consider indecomposable representations that are not necessarily irreducible. The next problem is how to control these theories that do not satisfy this requirement of traditional rational CFTs. Many of the papers cited above try to do so by considering extended chiral algebras [21], such that the number of indecomposable representations is finite, so that in this sense one may speak of “rational logarithmic CFTs”. But many applications, such as to disordered systems, and all the theories we consider in this paper, appear to possess only the Virasoro algebra as the chiral algebra, and infinitely many (non-isomorphic) indecomposable representations appear as local fields. In addition, these theories may even fail to be quasirational, that is infinite sums of local fields may be produced in ope’s. In these irrational CFTs, it is not clear how the analysis is to be organized.

We propose that the use of symmetry algebras that commute with the *full* chiral algebra [22] can be a powerful organizing principle, even when semisimplicity of the algebraic structures is lost. These symmetries and structures can be usefully studied by starting with lattice models. The idea of gaining understanding of logarithmic CFT (LCFT) by studying lattice realizations in details has been around for a while, and put forward most recently (albeit without the algebraic connotations we consider crucial) in [23]. It does open a fruitful new route to progress. Indeed, the lattice equivalent of

the reducible but indecomposable Virasoro representations in LCFT is the non semi-simplicity of the various algebras underlying the lattice models. [Examples include the Temperley-Lieb (TL) algebra or $U_q(\mathfrak{sl}_2)$, neither of which is semi-simple when q is a root of unity; such values are usually those of greatest interest.] A great deal of progress has been made in the mathematics literature in this area over the last fifteen years or so. Our strategy will be to exploit this progress and infer from it results about the LCFTs.

More precisely, the lattice models we deal with all have a similar structure (we discuss here only the boundary case, even though many results generalize to the case of the bulk). The local Hamiltonian densities generate an associative algebra (in our examples, the TL algebra) whose commutant, in the lattice Hilbert space of interest, is $U_q(\mathfrak{sl}_2)$ in the example of the spin-1/2 chain, or in the supersymmetric models in Refs. [20, 22] a remarkably large algebra whose representations are in “Morita equivalence” with those of $U_q(\mathfrak{sl}_2)$. Using results from the mathematics literature [24, 25, 26, 27] and others we derive [22], we can analyze in full detail the structure of these two commuting algebras (which are in so-called Ringel duality) and their representations in the lattice model. It is then a natural step to conjecture the corresponding structure in the continuum limit: the algebra of Hamiltonian densities goes over to the Virasoro algebra [28, 29], while the commutant goes over to an algebra of non-local charges. The general structure of the theory is one in which the Virasoro algebra commutes with a certain symmetry algebra, but the latter is *not* any kind of current algebra, and nor is it a Yangian. The structure of indecomposable representations in the lattice model goes over to the continuum limit, and we thus obtain information about the representations of the Virasoro algebra in our logarithmic theories. These representations are typically reducible but indecomposable with a diamond (or quartet) shape and made out of two “standard modules” connected by some “glue”. They bear a lot of resemblance to the so-called staggered modules introduced abstractly in [30]. The full structure of the Hilbert space involves big indecomposable representations of Virasoro and of its commutant, and can be exhibited in the form of what we will call a “staircase diagram”.

Although our approach does not so far yield much information on the detailed structure of the Virasoro indecomposable representations, it is enough to give access to the fusion rules for our logarithmic theories. We work these out in detail in some cases.

The remainder of this paper is organized as follows. Section 2 describes the general framework of the lattice models, algebraic structures, and the continuum limit, for q not a root of unity, and also the general features of cases in which q is a root of unity. The following three sections go into details about particular examples of special interest. Each one begins with a summary of the physical systems considered, viewed as loop models, or as field theories of certain nonlinear sigma models or Landau-Ginzburg theories, then describes the algebraic structures leading to the continuum limit, and ending with the fusion rules. Section 3 is the case $q = i$ (or central charge $c = -2$ in the continuum limit), which corresponds to symplectic fermions or dense polymers. Section 4 is the case $q = e^{i\pi/3}$ ($c = 0$ in the continuum limit), which corresponds to percolation, the spin quantum Hall transition, or a certain nonlinear sigma model at $\theta = \pi$. Section 5 is the case of dilute polymers ($c = 0$), or of the critical point in a certain nonlinear sigma model or Landau-Ginzburg theory. For this theory we discuss the continuum directly, and not the lattice models. Finally, section 6 is a more physical discussion of the structures for the cases with $c = 0$, using operator product expansions, and making contact with the work of Gurarie and Ludwig [17, 18].

To make reading easier, we provide a list of notations for some of the algebraic objects we introduce, all of which implicitly depend on q or the central charge c , and also on length L for the TL algebra:

- T_j : tilting modules over $U_q(\mathfrak{sl}_2)$
- P_j : projective modules over $TL_{2L}(q)$
- r_j : standard modules over the Virasoro algebra
- R_j : simple (or irreducible) modules over the Virasoro algebra
- $\mathcal{R}_j$: projective modules over the Virasoro algebra

2 Algebraic structure on the lattice and in the continuum

What we will do here is (i) take the continuum limit of the lattice model on a strip to obtain the representation content of the states, and (ii) also use the lattice to infer fusion rules for the boundary CFT.

In the companion paper [22], we showed that the commutant algebras of the Temperley-Lieb algebra in some supersymmetric lattice models are Morita equivalent [31] to $U_q(\mathfrak{sl}_2)$. Accordingly, here we can use the $U_q(\mathfrak{sl}_2)$ -invariant spin-1/2 chain, secure in the knowledge that the symmetry structure is the same in the supersymmetric constructions that apply when $m = q + q^{-1}$ is an integer. We often refer to representations as modules, as these are essentially the same things for our purposes (however, in the more mathematical discussions, saying “representation” in place of “module” would be quite misleading in some places, especially when preceded by “projective”). Then an irreducible representation is the same thing as a simple module, and it is also common to speak of modules being “over” the algebra that they represent. We assume familiarity with basic algebraic concepts such as sub- and quotient-objects, here usually for modules. For a general reference on the algebraic background, we suggest Ref. [31].

2.1 XXZ spin-1/2 chain, TL algebra and continuum limit: generic case

The Hamiltonian of the $U_q(\mathfrak{sl}_2)$ -invariant spin-1/2 chain is essentially the XXZ chain, but with some boundary terms (here the σ operators are Pauli matrices):

$$H = \frac{1}{2} \sum_{i=0}^{n-2} \left(\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \frac{q + q^{-1}}{2} \sigma_i^z \sigma_{i+1}^z \right) + \frac{q - q^{-1}}{2} (\sigma_0^z - \sigma_{n-1}^z) \quad (2.1)$$

The Hamiltonian H is built from generators e_i ,

$$e_i = \frac{q + q^{-1}}{4} - \frac{1}{2} (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) - \frac{q + q^{-1}}{4} \sigma_i^z \sigma_{i+1}^z - \frac{q - q^{-1}}{4} (\sigma_i^z - \sigma_{i+1}^z) \quad (2.2)$$

of the TL algebra $\text{TL}_n(q)$, which is defined as the algebra generated by a set of elements e_i , $i = 0, 1, \dots, n-2$, subject to the relations

$$e_i^2 = m e_i, \quad (2.3)$$

$$e_i e_{i\pm 1} e_i = e_i, \quad (2.4)$$

$$e_i e_{i'} = e_{i'} e_i \quad (2.5)$$

and no other relations algebraically independent of these. These are satisfied by the above expressions in the spin-1/2 chain. Then $H = -\sum_{i=0}^{n-2} e_i$ (up to an irrelevant additional constant). All future references to the spin-1/2 or XXZ chain are to this model.

First we briefly summarize the case of q generic (i.e., q not a root of unity) and its continuum limit. For q not a root of unity, the TL algebra is semisimple and has irreducible representations (all such will be referred to as simple modules hereafter) labeled by $j = 0, 1/2, 1, \dots, n/2$ but restricted to values such that $n/2 + j$ is an integer. The dimensions of these simple modules are d_j ,

$$d_j = \binom{n}{n/2 + j} - \binom{n}{n/2 + j + 1}. \quad (2.6)$$

These modules are referred to as standard (or Specht) modules.

The commutant of the TL algebra in the spin-1/2 chain is a finite-dimensional homomorphic image of $U_q(\mathfrak{sl}_2)$, that we will denote $U_q(\mathfrak{sl}_2)^{(n)}$ (it is also called the q -Schur algebra). For q generic, its simple modules are again labeled $j = 0, 1/2, 1, \dots, L/2$, where $n/2 + j$ is an integer. Throughout the paper, we will use the familiar term “spin” to refer to j , unless otherwise noted. These modules have dimensions $D_j = 2j + 1$, independent of n except that $j \leq n/2$. They are called Weyl modules [or standard modules of $U_q(\mathfrak{sl}_2)$] (we can drop the superscript n as they are independent of it, and the

limit $n \rightarrow \infty$ can be taken for these). They are the deformations to $q \neq 1$ of the usual irreducibles of $SU(2)$.

If two chains of lengths n_1, n_2 are joined end to end (with n_1 on the left), one obtains fusion rules for both algebras. The states of a single chain of length n_1 fall into multiplets (simple modules) of the tensor product of commuting algebras, $TL_{n_1}(q) \otimes U_q(\mathfrak{sl}_2)^{(n_1)}$ which are labeled by j ; these modules are the tensor product of the corresponding simple modules of the two respective algebras. When the two chains are joined, the resulting chain of length $n = n_1 + n_2$ can be analyzed in the same way. If the standard modules of $U_q(\mathfrak{sl}_2)$ are denoted V_j , then tensor products of such modules (where the tensor product is defined as in the usual way for vector spaces) decompose fully into a direct sum,

$$V_{j_1} \otimes V_{j_2} = \bigoplus_{j=|j_1-j_2|}^{j_1+j_2} V_j, \quad (2.7)$$

exactly as for $SU(2)$. The standard modules of TL must behave in a similar way, but now the product operation is defined using induction (which may be familiar from induced representations in group representation theory) from the product space as a module over $TL_{n_1}(q) \otimes TL_{n_2}(q)$ to a module over $TL_n(q)$, which contains the former as a subalgebra. The general theory described in Ref. [22] implies that indeed the “fusion rule” for this definition of a product of TL modules must take the same form as that above for $U_q(\mathfrak{sl}_2)$ (note this product is a functor, and not just a decomposition of the tensor product of vectors; in particular the dimensions of the tensor product of the modules on the left and the sum of those on the right are not equal).

Now we pass to the continuum limit. From this point on, we consider only $n = 2L$ even for the time being. (This does no harm as the modules with j integer close on themselves under products, as we see in the fusion rules above, and will see for non-generic q below. This restriction corresponds to the “oriented loops” models in ref. [22].) In the continuum limit we consider only the Fourier modes of the e_i s of wavevectors k with kL fixed, and also consider only states spanned by eigenstates with energies E (measured from the ground state energy) with EL fixed. Then the Fourier modes of the e_i s become the Virasoro generators L_n , $n = \dots, -2, -1, 0, +1, \dots$, [28]. The standard modules of the TL algebra, labeled by j , become certain modules of the Virasoro algebra, which we will also call standard, and which are still labeled by j . These modules are highest weight modules, that is they are generated by the action of the L_n s on the state of lowest energy or of lowest L_0 eigenvalue in the module. They are not, however, Verma modules, which are constructed from a highest weight state by lowering by L_n s with $n < 0$, and using only the commutation relations of the Virasoro algebra. The conformal weight h of the module, which is the L_0 eigenvalue of the highest weight state, takes one of the values $h_{r,s}$ at which a null vector (the highest weight in a submodule) appears in the Verma module, which must be set to zero in order to obtain a simple module. For generic values of the central charge, there is only a single null vector, and only when the conformal weight takes one of the values in the Kac table [32, 33]

$$h_{r,s} = \frac{[(x+1)r - xs]^2 - 1}{4x(x+1)}, \quad (2.8)$$

in which x is determined by the central charge

$$c = 1 - \frac{6}{x(x+1)}. \quad (2.9)$$

The generic values of c are those where x is irrational. When we take the limit of spin-1/2 chain, the resulting central charge follows from $q = e^{i\pi/(x+1)}$. The conformal weights that occur are $h_{1,1+2j}$ in the Kac table. In the corresponding Verma module, the null vector is at conformal weight $h_{1,-1-2j}$. Hence the Virasoro character of the simple modules that appear as direct summands in the $L \rightarrow \infty$ limit of the chain are (the trace is over states in the simple module only)

$$\text{Tr } \hat{q}^{L_0} = \frac{\hat{q}^{h_{1,1+2j}} - \hat{q}^{h_{1,-1-2j}}}{P(\hat{q})}, \quad (2.10)$$

where $\hat{q}$, $|\hat{q}| < 1$ is a parameter, $\hat{q} = e^{-\pi\beta/2L}$, where β is the inverse temperature (the length in the imaginary time direction), and $P(\hat{q})$ is the inverse of the Euler partition function, and is related to the Dedekind η function,

$$P(\hat{q}) = \prod_{r=1}^{\infty} (1 - \hat{q}^r) = \hat{q}^{-1/24} \eta(\hat{q}). \quad (2.11)$$

Thus the partition function of the continuum limit of the XXZ chain in the generic case is

$$\lim \text{Tr} e^{-\beta(H-E_0(L))} = \text{Tr} \hat{q}^{L_0} = \sum_{j=0}^{\infty} (2j+1) \frac{\hat{q}^{h_{1,1+2j}} - \hat{q}^{h_{1,-1-2j}}}{P(\hat{q})}. \quad (2.12)$$

Here the continuum limit is taken with $\hat{q}$ fixed, after subtracting the ground state energy $E_0(L)$ of the finite- L chain from the Hamiltonian, and the trace is over all the states of the spin-1/2 chain. [We emphasize that this is the ordinary trace on the vector space, not a q -trace; it corresponds to what were called the modified partition functions in Ref. [20].] If instead the subtraction were the length times the ground state energy density $e_0(L) = \lim_{L \rightarrow \infty} E_0(L)/L$ of the chain as $L \rightarrow \infty$, this expression would be multiplied by $\hat{q}^{-c/24}$. This is the way we will define the partition function here and in the following:

$$Z = \lim \text{Tr} e^{-\beta(H-e_0(L)L)} = \text{Tr} \hat{q}^{L_0 - c/24} = \sum_{j=0}^{\infty} (2j+1) \frac{\hat{q}^{h_{1,1+2j} - c/24} - \hat{q}^{h_{1,-1-2j} - c/24}}{P(\hat{q})}. \quad (2.13)$$

Let us note that this partition function can also be obtained by field-theoretic (rather than algebraic) means, by bosonizing the spin-1/2 chain or the 6-vertex model, that is, by representing it by a scalar field with a background charge, and taking the continuum limit [34, 35].

It is known that [33], as a consequence of the null vectors that were set to zero, the operator products of the fields $\phi_j(z)$ that correspond to the states in the modules decompose according to the fusion rules

$$\phi_{j_1} \times \phi_{j_2} = \sum_{j=|j_1-j_2|}^{j_1+j_2} \phi_j. \quad (2.14)$$

These clearly correspond to those we inferred from the induction product in the chains, or from the $U_q(\mathfrak{sl}_2)$ symmetry of the whole construction. In the following we use similar reasoning to infer fusion rules also in the non-generic cases in which representations are not fully decomposable.

In this procedure, we have constructed the fusion product algebraically by joining two chains end to end and taking the continuum limit. One may wonder whether this really yields the operator product of corresponding local boundary fields in the CFT, which is usually pictured as bring points on the boundary together. We will not discuss this fully, but provide some heuristic arguments. If we view joining the chains as an event in time, then it corresponds to a picture like that shown in Fig. 1, in which the continuum limit has been assumed. (The chains are shown as equal in length, though more generally the ratio of their lengths can be arbitrary, but is held fixed when the continuum limit is taken.) If we wish to consider the fusion of states that are L_0 eigenstates, for example, then the two initial states may be prepared at early imaginary time $t \rightarrow -\infty$, and the matrix element with a desired final state can be taken, by preparing that state at $t \rightarrow +\infty$. As conformal transformations move points within the interior of the domain (and even deform the domain), it is natural that the tensor product of two copies of Virasoro merges into a single copy of Virasoro, as a proper subalgebra. This occurs because the continuum analog of the extra generator e_{n_1-1} must be included along with the others. Then the corresponding actions of these algebras on the states are related by the induction functor. Thus the use of the induction functor for Virasoro is natural in the slit-strip picture (i.e., in the operator formalism). The conventional picture of the incoming and outgoing states as corresponding to local fields at a point on the boundary, which are brought together in an ope, is recovered by using a conformal mapping from the slit-strip to the unit disk, or the upper half plane, with the two incoming and one outgoing state(s) at time $\rightarrow \pm\infty$ represented by points on the boundary. This conformal mapping is described for example in Ref. [36]. Incidentally, it has often been remarked in the CFT




Figure 1: The “slit-strip” picture of open string interactions, or of joining two chains end to end, with time increasing upwards. The shading indicates the interior of the domain.

literature that the fusion product is not simply a tensor product of representations of Virasoro (or more generally, the chiral algebra). The fusion product defined as the induction functor for joining strips end to end should agree with the “deformed tensor product” defined in some of these references [21], at least for the present case of boundary CFT.

The states or fields we find in this construction based on the TL algebra at generic q correspond to the first column of the Kac table. The first row of the Kac table has identical fusion rules, and so one would expect a lattice construction to lead to this theory also. Such a model is the $O(m)$ loop model [37, 38, 39] and its supersymmetric versions [20, 22] in which the critical theory (which exists for $-2 < m \leq 2$) is referred to as the dilute theory, and where another algebra (essentially a two colour version of the TL algebra) replaces the TL algebra. We emphasize that it has the same symmetry as the TL (dense loops) model, and that it too comes in two versions, oriented loops (only integer $j \geq 0$) and unoriented loops (including all integer values of $2j \geq 0$).

2.2 Non-generic cases

When q is a root of unity (but $q \neq \pm 1$), we will define $r > 1$ to be the smallest integer such that $q^r = \pm 1$. The structure of both TL and $U_q(\mathfrak{sl}_2)$ changes, and this can be traced to the vanishing of the q -integers $[N]_q = (q^N - q^{-N})/(q - q^{-1})$ for some N for such q values: $[N]_q = 0$ for $N \equiv 0 \pmod{r}$ [40, 25, 26].

For q a root of unity, standard modules of both TL and $U_q(\mathfrak{sl}_2)$ can be defined, and have the same dimensions d_j , $2j + 1$, respectively, as before. However, they are not all irreducible. It turns out that for j such that $[2j + 1]_q = 0$, they remain irreducible, while for other j , they contain a proper submodule but are indecomposable. For both algebras, the reducible standard modules have the same structure: the largest proper submodule (or radical) of the module is irreducible, and the quotient by this submodule is also. Then we will represent the structure of such a module by a diagram like

$$\begin{array}{c} \circ \\ \searrow \\ \bullet \end{array} \quad (2.15)$$

Here the open circle represents the states in the simple quotient module or “top” (or “head”), and the

closed circle the simple submodule or “foot”. The arrow represents the action of the algebra; there is some element of the algebra that maps the top to the foot, but not vice versa, as well as elements that map the top into itself and the foot into itself. The positions of the circles on the page is not particularly important, as the key information is the arrows and (for these standard modules) whether a circle is open or closed. Below, we will also represent the structure of other modules by similar, though more complex, diagrams. In these, each circle is a nonzero simple subquotient module, and arrows show the action of the algebra other than within the simple subquotients, with the convention that composites of arrows should also be understood as present implicitly. In our examples, a head and a foot can usually be defined uniquely, but there will be other subquotients in between.

Later, the positions of the circles will be replaced by additional information such as the spin or dimension of the simple module in question. The general rule for the spin values can be given here: For either $U_q(\mathfrak{sl}_2)$ or TL, we label standard modules by $j = 0, 1, \dots, L$. We choose the labels for the simple modules to be $j = 0, 1, \dots, L$ also (or in some cases, a subset of these). For reducible standard modules of $U_q(\mathfrak{sl}_2)$, the simple module at the *top* has the same value of j ; the foot is a simple module isomorphic to a simple module with *smaller* j . For the TL algebra, it is the same, except that the foot has a larger j than the head.

When the XXZ chain is analyzed for q a root of unity, it decomposes under the TL algebra into a direct sum of TL modules. When analyzed under $U_q(\mathfrak{sl}_2)$, it likewise is a direct sum. However, in neither case are the direct summands the standard modules just described, except for those standard modules that remain simple at this q , some copies of which appear as in the generic cases. For the reducible standard modules, each algebra action mixes them, mapping states from one standard to another [40, 24, 25, 26]. The direct summand modules will be referred to as “tilting” modules, in the case of either algebra. (We should point out that non-semisimple algebras typically have infinitely many distinct, i.e. mutually non-isomorphic, indecomposable modules, even when the algebra is finite dimensional. It becomes important to explain *which* particular types of indecomposable modules are of interest.) The tilting modules turn out to have a fairly simple structure also, consisting of two standard modules with arrows connecting them, typically in the form:

Thus, the module is seen to contain a submodule that is a standard module, consisting of an open and a closed circle together with an explicit arrow that connects them. The quotient by this submodule is also standard. The detailed structure of all these modules will be exhibited for several important values of q (or m) in the following sections, as will the inter-relation of the commuting TL and $U_q(\mathfrak{sl}_2)$ actions.

In addition to the tilting modules, other important modules are the projective modules. Projective modules may be defined as occurring as a direct summand in some free module, or in several other equivalent ways. We will be especially interested in the indecomposable projective modules of the TL algebra. For finite dimensional algebras, there is one indecomposable projective module (up to isomorphism) corresponding to each distinct simple module; the latter appears as the top (the unique simple quotient module) of this projective module. This fact makes it fairly straightforward to read off the indecomposable projective modules from the structure of the XXZ chain as a direct sum, since the algebra acts faithfully. In many cases, the tilting modules are also indecomposable projective modules.

Joining chains of lengths L_1, L_2 end to end leads to natural operations making a tensor product of modules over $U_q(\mathfrak{sl}_2)$ into another such module. These agree with the product operations usually defined for $U_q(\mathfrak{sl}_2)$ as a quantum group. This product decomposes as a direct sum of indecomposable modules, and as tilting modules are defined as direct summands in the XXZ chain, the tensor product of tilting modules is again tilting. The fusion rules for the tilting modules are then those of most basic interest (and are determined by the decomposition of a product of indecomposable tilting modules into indecomposables). Also, as this product is just the ordinary tensor product operation of vector spaces, the sum of the dimensions of the indecomposables produced is the product of the dimensions of the

modules we began with. For q a root of unity, the fusion rules are however modified from those in the generic cases, and will be discussed in the examples below.

In our previous paper [22], we also explained how a product is defined on the TL modules when joined end to end, based on induction as mentioned earlier. In this case, the dimensions of modules are not conserved, but there are again fusion rules. Crucially, the module induced from a tensor product of projective modules of the two short chains is a projective module of the chain of length $L_1 + L_2$, and can be decomposed as a direct sum of indecomposable projective modules. Hence we can define fusion rules for products of projective modules, which are again determined by the decomposition of a product of indecomposable projective modules into indecomposable projective modules. Most importantly, the fusion rules for indecomposable projective modules of TL are the *same* as those for indecomposable tilting modules of $U_q(\mathfrak{sl}_2)$ [22].

We also discussed previously [22] how the limit $L \rightarrow \infty$ can be taken, both at a purely algebraic level, and focussing on the low-energy states as a continuum limit. The first can be well-defined mathematically, while the second is presently less rigorous and more heuristic. For the symmetry structure, given by $U_q(\mathfrak{sl}_2)$ or a Morita equivalent of it, the purely algebraic construction is sufficient, as it yields a limit algebra, which is $U_q(\mathfrak{sl}_2)$ for the XXZ case. The dimensions of modules are stable as L increases, but additional modules appear. The fusion rules are also given by the finite-dimensional analysis and remain the same as L increases [and in discussing $U_q(\mathfrak{sl}_2)$ above, it was natural to ignore the distinction between it and its finite-dimensional quotients]. On the other hand, for the TL algebra, the dimensions of corresponding modules increase with L , and there seems to be no purely algebraic way to define the limiting algebra that is suitable for our purposes [22]. Nonetheless, the fusion rules behave stably (as they are controlled by the symmetry). In the purely algebraic limit, the limit of a projective module is still a projective object in the limiting category of modules. In the continuum limit, all evidence suggests that the modules of TL become Virasoro modules. The limits of projective modules will still be referred to as projective, and continue to play an important role. The standard (resp., tilting) modules of TL become standard (resp., tilting) modules of Virasoro, also. We note that the Virasoro algebra may not act faithfully on the limiting space of states. This is to be expected as the representations that occur as summands are quite restricted, so that the algebra appears smaller—it is a quotient of Virasoro. We can see this already for generic q , where the standard modules became the irreducible representations of Virasoro that lie in the first column of the Kac table. It is possible that the modules we call projective do obey the general definition of projective modules over this smaller algebra, even if they do not for the Virasoro algebra itself. Also, there could in principle be elements of the algebra not in the Virasoro algebra (though this seems unlikely in the spin-1/2 chains and related theories studied here).

When q is a root of unity, the corresponding central charge is one of the special values of Kac and BPZ (in which x above is rational), at which the standard modules may acquire additional null vectors. When this occurs, the standard module of Virasoro now has the same submodule structure as the standard TL module illustrated above. The positions in the first column of Kac table at which this occurs agree exactly with what we would expect based on the TL algebra. Similarly, we will infer that the projective and tilting Virasoro modules relevant to the analysis of the XXZ chain (to operators as well as states, by the state-operator correspondence) have structures like that above (when reducible). Finally, the fusion rules must be those inferred from finite size systems, and for the projective modules are again the same as for finite size L . These can in turn be obtained from those on the $U_q(\mathfrak{sl}_2)$ side, which is very convenient as one can use characters there. This illustrates how symmetry becomes useful in analyzing the structure of a CFT.

Thus, in the following sections, the ideas of this part will be worked out in several instances of physical or geometric interest.

3 The case $m = 0$ ($c = -2$) — dense polymers

In this section, we consider the case of the open XXZ chain of even length $2L$, with $m = 0$, so $q = i$ and $r = 2$. When the partition function is expanded as a sum over loops, this model assigns a factor $m = 0$ to each loop, so describes dense polymers. This case also corresponds (by Morita equivalence of

symmetry algebras) to the $\mathfrak{gl}(n|n)$ supersymmetric chains, and to nonlinear sigma models with target space $\mathbf{CP}^{n-1|n}$ at a particular value of the coupling constant, as described in Ref. [20]. For $n = 1$, the $\mathfrak{gl}(1|1)$ spin chain actually coincides with the $U_q(\mathfrak{sl}_2)$ or spin-1/2 chain, and both are free fermion systems. In the continuum limit, the latter theory is the free symplectic fermion theory.

The decomposition of the space of states into a sum of indecomposable representations (tilting modules) under $U_q(\mathfrak{sl}_2)$ has been well studied. It turns out that none of the indecomposable summands are simple; all are reducible. Following the notation described in the previous section, each tilting module T_j has the structure corresponding exactly to that in diagram (2.16) [40, 25, 26]:

$$T_j : \begin{array}{ccc} & j-1 & \\ \swarrow & & \searrow \\ j-2 & & j \\ \searrow & & \swarrow \\ & j-1 & \end{array} , \quad j = 1, \dots, L.$$

The nodes correspond to simple modules and are labeled by spin values; the simple module of spin $j' \geq 0$ has dimension $j' + 1$. (Nodes with negative spin values are viewed as zero dimensional and can be omitted.) For some purposes, the module T_0 for $j = 0$ can also be viewed as a tilting module of $U_q(\mathfrak{sl}_2)$ but does not occur in the chain because its multiplicity is zero (this is special to the $m = 0$ case); in the present section we will view it also as zero. Each pair of nodes that can be connected by a southwest arrow forms a standard module, as in diagram (2.16). As advertised, the dimension of the standard module of spin j' is $2j' + 1$; conversely, this together with a knowledge of the structure enables one to calculate recursively the dimensions of the simple modules. The total dimension of the module is thus $4j$. We emphasize that the same structure applies to the corresponding tilting modules for the enlarged symmetry algebra $\mathcal{A}_{n|n}(2L)$, by Morita equivalence, though the dimensions are different. Similar statements will be true for the symmetry algebras throughout, but will not usually be made explicit.

For $L = 3$ (i.e. six spins 1/2) for example, the j values of the indecomposable summands T_j are $j = 1, 2, 3$. The full decomposition of the chain under $U_q(\mathfrak{sl}_2)$ (where $\times n$ denotes multiplicities of the indecomposable tilting modules) is

$$\begin{array}{ccc} 0 & & 1 & & 2 \\ & \searrow & \swarrow \quad \searrow & & \swarrow \quad \searrow \\ & 1 & 0 & 2 & 1 & 3 \\ & \swarrow & \nwarrow \quad \swarrow & & \nwarrow \quad \swarrow \\ 0 & & 1 & & 1 \end{array} \times 5 , \quad \times 4 , \quad \times 1 \quad (3.1)$$

and the check on the dimensions is

$$(1 + 2 + 1) \times 5 + (2 + 1 + 3 + 2) \times 4 + (3 + 2 + 3 + 4) \times 1 = 64. \quad (3.2)$$

Knowing the decomposition under (an image of) $U_q(\mathfrak{sl}_2)$, one can also find directly the structure of the modules of the commutant algebra of $U_q(\mathfrak{sl}_2)$ in the spin-1/2 chain. This by definition consists of all endomorphisms (linear maps of the vector space into itself) that commute with the $U_q(\mathfrak{sl}_2)$ action. When the algebra is semisimple, as for $U_q(\mathfrak{sl}_2)$ at generic q , such endomorphisms can only map a copy of a simple module to another copy of the same one, and so the commutant is also semisimple, and has simple modules of dimensions equal to the multiplicities of the simple modules of the given algebra (this was used in Ref. [22]). For non-semisimple algebras, as here, there are such endomorphisms that map the tilting modules to themselves and which combine to form full matrix algebras. Indeed, the multiplicities of the $U_q(\mathfrak{sl}_2)$ tilting modules are the dimensions of the simple modules of its commutant. But there are also endomorphisms that map the indecomposable tilting modules into one another. This can occur when a tilting module has one or more subquotient modules that also occur in another tilting module. In fact the “top” of the indecomposable tilting module, which is a source but not a sink for arrows, must map onto a non-top part of the other tilting module, and all subquotients that lie below the top must also map such that the arrows are preserved. In most cases, the foot of the




Figure 2: The structure of the spin-1/2 chain for $q = i$ and $2L = 6$ sites, as a representation of $U_q(\mathfrak{sl}_2) \otimes \text{TL}_{2L}(q)$, or the same for the $\mathfrak{gl}(n|n)$ -supersymmetric spin chain with $2L = 6$ as a representation of $\mathcal{A}_{n|n}(2L) \otimes \text{TL}_{2L}(q)$.

tilting module, and some subquotients above it, are mapped to zero. Thus the endomorphisms of the space of the spin-1/2 chain can be read off from the structure and multiplicities of the indecomposable tilting modules. For the TL representation in the spin-1/2 chain, the commutant algebra of $U_q(\mathfrak{sl}_2)$ obtained in this way is isomorphic to the TL algebra; the two algebras form a “dual pair”. (If instead the TL structure were given, one could follow the same procedure to find the structure of the modules of its commutant.)

In the present case, the indecomposable tilting modules T_j have non-trivial homomorphisms into neighboring tilting modules $T_{j \pm 1}$ only. There are also (for $j > 0$) non-trivial homomorphisms of T_j into itself, which map the head to the foot, which have the same j value. The commuting actions of $U_q(\mathfrak{sl}_2)$ and its commutant $\text{TL}_{2L}(q)$ can be represented in a “staircase” diagram that generalizes those for $U_q(\mathfrak{sl}_2)$ or TL separately, shown for $L = 3$ in Fig. 2. This shows the action of both algebras, and thus the tensor product $U_q(\mathfrak{sl}_2) \otimes \text{TL}_{2L}(q)$, on the states of the chain. [The diagram is the same if we consider $\mathcal{A}_{n|n}(2L) \otimes \text{TL}_{2L}(q)$ for the supersymmetric chains.] The circles at Cartesian coordinates (j, j') represent simple subquotient modules over this product algebra. $U_q(\mathfrak{sl}_2)$ acts by the horizontal arrows (that is, its action preserves j'), while $\text{TL}_{2L}(q)$ acts “vertically” (that is, preserving j), and these actions commute (readers are invited to check that this is consistent, because of the arrangement of the arrows as “commuting squares”). Once again, composites of arrows are arrows that are implicitly present. Each circle is a subquotient module with a multiplicity of one when viewed as a $U_q(\mathfrak{sl}_2) \otimes \text{TL}_{2L}(q)$ module; some values (j, j') do occur twice, and those circles have been separated slightly for clarity. The indecomposable tilting modules of $U_q(\mathfrak{sl}_2)$ can be recovered by ignoring vertical arrows, and stretching out the components of the diagram to recover the diagrams above. Likewise, the indecomposable tilting modules of the TL algebra are obtained by ignoring the horizontal arrows. (If we had started knowing those, then we could have obtained the tilting modules for the commutant of TL by a similar procedure as we described.) We now turn to the structure of these in some examples. For this $m = 0$, L integer case, the general form is always a single indecomposable module under the product algebra, and for $L > 3$ simply extends up to larger j values.

In detail for $L = 3$, the TL indecomposable tilting modules (direct summands) have the following structure, in which the nodes are labelled by the dimension of the corresponding simple subquotient

module over TL, and arrows now denote action of the TL algebra [24, 25, 26]:

$$\begin{array}{c} (5) \\ \searrow \\ (4) \\ \swarrow \\ (5) \end{array} \times 1, \quad \begin{array}{c} (4) \\ \swarrow \quad \searrow \\ (1) \quad (5) \\ \swarrow \quad \searrow \\ (4) \end{array} \times 2, \quad \begin{array}{c} (1) \\ \swarrow \quad \searrow \\ (4) \quad (1) \\ \searrow \\ (1) \end{array} \times 3, \quad (1) \times 4, \quad (3.3)$$

with total dimension

$$(5 + 4 + 5) \times 1 + (5 + 4 + 4 + 1) \times 2 + (1 + 4 + 1) \times 3 + 1 \times 4 = 64 \quad (3.4)$$

Notice here the appearance of multiplicities from the $U_q(\mathfrak{sl}_2)$ analysis as dimensions of simple TL modules, and vice versa. Using the labels j for the TL simple modules as already described, eq. (3.3) becomes

$$\begin{array}{c} 1 \\ \searrow \\ 2 \\ \swarrow \\ 1 \end{array} \times 1, \quad \begin{array}{c} 2 \\ \swarrow \quad \searrow \\ 1 \quad 3 \\ \swarrow \quad \searrow \\ 2 \end{array} \times 2, \quad \begin{array}{c} 3 \\ \swarrow \quad \searrow \\ 2 \quad 3 \\ \searrow \\ 3 \end{array} \times 3, \quad 3 \times 4, \quad (3.5)$$

For general $L > 1$, the pattern is similar, starting with a “trio” at $j = 1$, followed by quartets up to $j = L$, and then a single simple module also at $j = L$ (here we assign j to the TL tilting modules by the j of the head).

The tilting modules for TL can also be compared with the standard modules. The standard module labeled j for $\text{TL}_{2L}(q = i)$ consists of the simple modules $j, j + 1$, together with the TL action that connects them by an arrow, except for $j = L$ when the simple module is standard and one dimensional ($d_L = 1$), and $j = 0$ where the standard module is simple. For six spins 1/2 again, one has $d_0 = 5$, $d_1 = 9$, $d_2 = 5$, $d_3 = 1$. The indecomposable tilting modules consist of pairs of standard modules “glued” together by the action of some TL elements, similarly to $U_q(\mathfrak{sl}_2)$.

When the structure of the tilting modules is known as in (3.5), one can recursively calculate the dimensions d_j^0 of the simple modules of TL, starting from $j = L$. For $L = 3$ this gives $d_3^0 = 1$, $d_2^0 = d_2 - d_3 = 4$, $d_1^0 = d_1 - d_2^0 = 5$, $d_0^0 = d_0 - d_1^0 = 0$. Usually there is a simple module of TL for $j = 0$, but here for $m = 0$ it vanishes. (This is related with the fact that the partition function of the chain vanishes exactly). For general even $2L$, one finds the formula for the dimensions of simple TL modules for the $q = i$ case,

$$d_j^0 = \sum_{j'=j,j+1,\dots,L} (-1)^{j-j'} d_{j'} \quad (3.6)$$

for $j = 0, 1, \dots, L$.

We may also consider $U_q(\mathfrak{sl}_2)^{(2L)}$ and analyze it viewed as a module over itself, acting on the left or on the right (the left and right regular representations), or on both sides (i.e. as a $U_q(\mathfrak{sl}_2)^{(2L)} \otimes U_q(\mathfrak{sl}_2)^{(2L)}$ module, as the actions on the two sides commute). From the last of these, we can then make a similar staircase diagram for $U_q(\mathfrak{sl}_2)^{(2L)}$ as a module over $U_q(\mathfrak{sl}_2)^{(2L)} \otimes U_q(\mathfrak{sl}_2)^{(2L)}$, from which the left or right structure can be determined. The cellular ideals we reviewed in Ref. [22] can be readily seen in such a picture, as they form sets of circles and arrows mapped into themselves by the left and right action of $U_q(\mathfrak{sl}_2)^{(2L)}$. This is also useful in calculating the total dimension of the algebra, which is equal to $\dim U_q(\mathfrak{sl}_2)^{(2L)} = \sum_{j=0}^L (2j+1)^2$, even though it is not semisimple. One can do the same for the TL algebra, and then $\dim \text{TL}_{2L}(q) = \sum_{j=0}^L d_j^2$ ($= 132$ for $L = 3$). Similarly, for the spin-1/2 chain itself, we can point out that the total dimension is $2^{2L} = \sum_{j=0}^L (2j+1)d_j$, again just as if the algebras were semisimple. Of course all of these expressions for the dimensions follow from the independence of the dimensions of the standard modules from q , so that the formulas from the generic case always hold. This is also true for the other non-generic values of q , some of which are considered later. For the models with Morita equivalent symmetry algebras, $2j+1$ should be replaced by the dimensions $D'_j = [2j+1]_{q'}$, where $q' + q'^{-1} = 2n$ [22], but again it is true that the formulas can be obtained “as

if” the standard modules were simple (similar formulas also hold for the q -dimensions [in the $U_q(\mathfrak{sl}_2)$ case] and the superdimensions [in the superalgebra cases [22]]).

We will give here the expressions for the (super-)dimensions of the simple modules of our algebras $\mathcal{A}_{n|n}$, $n = 1, 2, \dots$, which are Morita equivalent to $U_q(\mathfrak{sl}_2)$ with $q = i$ [22]. First we point out that the dimensions $j + 1$ of the j th simple module of $U_q(\mathfrak{sl}_2)$ can be deduced from the structure of the standard modules, and the dimension $2j + 1$ of the latter, much like those for TL. They are obtained from the relations $j + 1 = (2j + 1) - j$, with dimension 1 for the simple module with $j = 0$, which follow from the existence of simple sub- and quotient modules of the each standard module. For the algebras in the supersymmetric models, the standard modules have dimensions $D'_j = [2j + 1]_{q'}$, where $q' + q'^{-1} = 2n$ [22]. Then as the structure of submodules and quotients is the same as for $U_q(\mathfrak{sl}_2)$ because of the Morita equivalence [22], we have for the dimensions D_j^0 of the simple modules the corresponding recurrence $D_j^0 = D'_j - D'_{j-1}$, with $D'_0 = D_0^0 = 1$ obviously corresponding to a simple module. This is solved by

$$\begin{aligned} D_j^0 &= D'_j - D'_{j-1} + D'_{j-2} - \dots \\ &= q'^{2j} + q'^{2j-4} + \dots + q'^{-2j} \\ &= [j + 1]_{q'^2}. \end{aligned} \quad (3.7)$$

Similarly, we may look at the quantum dimensions of the modules; for the algebra $\mathcal{A}_{n|n}$ the quantum dimensions are just the superdimensions. For the j th standard module, the superdimensions were denoted D_j in Ref. [22], given by $D_j = [2j + 1]_q$, which is the same as the q -dimension for $U_q(\mathfrak{sl}_2)$ because of the Morita equivalence with $U_q(\mathfrak{sl}_2)$ as ribbon Hopf algebras (here for the algebras with integer j only) [22]. In the present case, $D_j = (-1)^j$. The same equality applies to dimensions of other modules such as the simple modules, and thus we can compute the quantum dimensions for $U_q(\mathfrak{sl}_2)$ ($q = i$) simples and $\mathcal{A}_{n|n}$ simples as $D_j^0 = [j + 1]_{q^2}$ similarly. For the tilting modules, the quantum dimensions are $D_j^+ = D_j + D_{j-1}$ ($j = 1, 2, \dots$), and so $D_j^+ = 0$ for all $j \geq 1$.

For a finite-dimensional algebra, the indecomposable summands in the left regular representation are the indecomposable projective (left) modules. Thus we can read off the indecomposable projective modules. For the TL algebra (for which these are of particular interest) for $L = 3$, these are the same modules as the three tilting modules other than the singlet. The singlet cannot be an indecomposable projective module as it occurs as a quotient module of an indecomposable tilting module, which is impossible for a projective module (one of the alternative definitions for a projective module is that whenever it is a quotient of some module, it is isomorphic to a direct summand in that other module). The other three indecomposable tilting modules are acceptable, and are projective as follows automatically if the TL algebra is defined as the commutant of $U_q(\mathfrak{sl}_2)$ in this chain (this is an aspect of the key lemma [41]; see also Ref. [22]). There are no other indecomposable projective modules in the present case (for the other cases $m \neq 0$, we will see that there are also projective modules that are not tilting). So the indecomposable projectives of the TL algebra are

$$P_j : \begin{array}{c} 1 \\ \swarrow \quad \searrow \\ 2 \\ \swarrow \quad \searrow \\ 1 \end{array} \times 1, \begin{array}{c} 2 \\ \swarrow \quad \searrow \\ 1 \quad 3 \\ \swarrow \quad \searrow \\ 2 \end{array} \times 2, \begin{array}{c} 3 \\ \swarrow \quad \searrow \\ 2 \quad 3 \\ \swarrow \quad \searrow \\ 3 \end{array} \times 3, \quad (3.8)$$

This pattern is repeated for general $L > 0$ (integer); the indecomposable projectives P_j of TL for $q = i$ are the indecomposable tilting modules, other than the singlet at $j = L$:

$$P_j : \begin{array}{c} j \\ \swarrow \quad \searrow \\ j-1 \quad j+1 \\ \swarrow \quad \searrow \\ j \end{array}, \quad j = 1, \dots, L, \quad (3.9)$$

where once again we understand in this case that the simple modules with $j < 1$ or $j > L$ are zero and can be omitted. (For $L = 1$, one does have a two-dimensional indecomposable P_1 , because of the implicit arrow from the head $j = 1$ to the foot $j = 1$.)




Figure 3: Space of states of the continuum $m = 0$ theory, showing the commuting actions of Virasoro and $U_q(\mathfrak{sl}_2)$ (or $\mathcal{A}_{n|n}$).

We now pass to the continuum limit. Our strategy will be to deduce as much as possible from the lattice, and compare with known results from the representation theory of the Virasoro algebra. First, we can consider the purely algebraic limit $L \rightarrow \infty$ as discussed in Ref. [22]. The staircase diagram then becomes infinite; see Fig. 3, and the limits of the dimensions d_j^0 of the simple TL modules are also infinite. The more physical way of taking $L \rightarrow \infty$, called the continuum limit [22], focusses on low-energy states and the long-wavelength Fourier components of the e_i s. We expect that the Virasoro algebra emerges from the TL algebra in this limit [28], and that the staircase diagram still applies to this. In addition we will use the known information (as for the case of generic q) of the values of the central charge and a set of scaling dimensions that are associated with the Virasoro modules (or the scaling fields of the theory). This is done for convenience; we expect that with some additional effort these may themselves be established purely on the grounds of consistency with the algebraic structure encoded in the staircase diagram, since the structure of the $U_q(\mathfrak{sl}_2)$ modules is known and is unchanged in the $L \rightarrow \infty$ limit.

On specializing the generic results, eqs. (2.9), (2.8) for the central charge and the Kac table to $q = i$, so $x = 1$, one finds that $c = -2$, and the Kac formula reads

$$h_{rs} = \frac{(2r - s)^2 - 1}{8}, \quad (3.10)$$

where r, s are positive integers. The values appearing in the spectrum are

$$h_{1,1+2j} = h_{j,1} = \frac{(2j - 1)^2 - 1}{8} = \frac{j(j - 1)}{2}, \quad (3.11)$$

where we used symmetries of the Kac formula to restrict to the fundamental domain, and again here $j \geq 0$ is integer.

As the limits of the dimensions of the modules over TL are infinite, it is convenient to use instead the graded dimensions or characters defined using a trace over the module of the exponential of the Hamiltonian, or L_0 Virasoro generator. For the standard modules, which we will denote $r_{r,s}$, these were given in Sec. 2, and for the cases of interest here are

$$K_{1,1+2j} = \text{Tr}_{r_{1,1+2j}} \hat{q}^{L_0 - c/24} = \frac{\hat{q}^{(2j-1)^2/8} - \hat{q}^{(2j+3)^2/8}}{\eta(\hat{q})}. \quad (3.12)$$

$$(1,1) \longrightarrow (2,1) \longrightarrow (3,1) \longrightarrow$$

Figure 4: Chain of Verma modules embeddings for $c = -2$.

We may point out here that the full partition function of the model in the continuum limit is obtained by specializing the result for generic q :

$$Z = \text{Tr} \hat{q}^{L_0 - c/24} = \sum_{j=0}^{\infty} (2j+1) \frac{\hat{q}^{(2j-1)^2/8} - \hat{q}^{(2j+3)^2/8}}{\eta(\hat{q})}, \quad (3.13)$$

which as for the total dimension of the finite length chains is still valid at $q = i$, even though the modules involved are no longer fully decomposable. This is similar to the result given for the case $c = 0$ in the appendix of Ref. [20]. This result may be obtained using field-theoretic methods [34], and provides a check on our algebraic deductions. It is perhaps more natural to calculate a partition function as a quantum trace instead of a trace, which corresponds to a boundary condition in the (periodic, imaginary) time direction that does not break the $U_q(\text{sl}_2)$ or $\mathcal{A}_{n|n}$ symmetry. In that case $2j+1$ is replaced by D_j , and the result is valid for the continuum limit of the supersymmetric models as well as that of the spin-1/2 chain [20]. In the present case $q = i$, these partition functions (called “unmodified” in Ref. [20]) vanish identically (this is also true on the lattice). This follows because we can collect all states into tilting modules of the symmetry algebra, and the quantum dimensions D_j^+ of these tilting modules vanish. For the modified partition functions of the supersymmetric models, $2j+1$ must be replaced by D_j' as for the finite- L chains.

In studying the representation theory for highest-weight modules of the Virasoro algebra, it is conventional to use Verma modules. The Verma module generated from the highest weight vector with weight $h_{1,1+2j}$ has, for generic central charge, a singular vector at level $1+2j$, with value of the conformal weight $h_{1,-1-2j}$. That is, there is a submodule which is also highest weight with that conformal weight, and vectors in the submodule are orthogonal to all those in the Verma module. At $c = -2$, another singular vector appears at the lower level $h_{j+1,1} = h_{j,1} + j$, while now $h_{1,-1-2j} = h_{j+2,1} = h_{j,1} + 1 + 2j$. These submodules in turn contain submodules. This leads to a picture of Verma modules embedded into one another in a sequence, as shown in Fig. (4).

The representations obtained by factoring out from the Verma module the submodule generated from the “generic” singular vector at level $1+2j$ are the standard modules $r_{1,1+2j}$. At $c = -2$ this representation is reducible; from Fig. 4 it contains a submodule that is simple, and its structure is described by a diagram of the form (2.15). It possesses an irreducible quotient $R_{j,1}$, with character $\chi_{j,1}$, and the character of the submodule in $r_{1,1+2j}$ is $\chi_{j+1,1}$. Hence we have the equality

$$K_{1,1+2j} = \chi_{j,1} + \chi_{j+1,1}. \quad (3.14)$$

The calculation of $\chi_{j,1}$ from the embedding diagram Fig. 4 is standard, and gives

$$\text{Tr}_{R_{j,1}} \hat{q}^{L_0 - c/24} = \chi_{j,1} = \frac{\hat{q}^{(2j-1)^2/8} - \hat{q}^{(2j+1)^2/8}}{\eta(\hat{q})}, \quad (3.15)$$

and note that $\chi_{0,1} = 0$ (the continuum equivalent of the absence of the $(0,0)$ node in Fig. 2).

This shows that the structure of the standard Virasoro modules at $c = -2$ parallels that of the finite- L TL modules of the same j . We now go a step further and describe the structure of Virasoro modules that parallel the indecomposable projective modules of TL; these will be called indecomposable projective modules of Virasoro (the terminology is subject to caveats already mentioned in the previous section). The modules we want are the continuum limits of the finite- L indecomposable projective TL modules. Calling these indecomposable projective modules $\mathcal{R}_j$, we expect thus in particular to have the characters

$$\text{Tr}_{\mathcal{R}_j} \hat{q}^{L_0 - c/24} = K_{1,1+2j} + K_{1,1+2(j-1)} = 2\chi_{j,1} + \chi_{j-1,1} + \chi_{j+1,1}. \quad (3.16)$$




Figure 5: The submodule structure of the indecomposable projective modules of the Virasoro algebra. The vectors are highest weights in the corresponding simple subquotient modules. Open dots mean the corresponding vectors are not Virasoro descendants of any other state within the representation.

The structure of these modules, in the sense of the submodules that appear, is as follows. All are described by the same diagrams as the corresponding projectives P_j of TL, and all except that for $j = 1$ are “quartets” since the effects at $j = L$ have disappeared as $L \rightarrow \infty$. Each circle in the diagram is a simple subquotient module, and for the Virasoro algebra these are highest weight modules. In more detail, we should first have an irreducible representation $R_{j,1}$ generated by a highest weight vector $V_{j,1}$ at the “foot”; then a standard submodule $r_{j-1,1}$ generated by a highest weight vector $v_{j-1,1}$, in which $V_{j,1}$ appears as a singular vector. Then a state $V'_{j,1}$ which forms a two dimensional Jordan cell (of L_0) with $V_{j,1}$:

$$\begin{aligned} L_0 V'_{j,1} &= h_{j,1} V'_{j,1} + V_{j,1}, \\ L_0 V_{j,1} &= h_{j,1} V_{j,1}. \end{aligned} \quad (3.17)$$

$V'_{j,1}$ is not a highest weight vector, and positive modes of Virasoro (i.e. L_n 's with $n > 0$) map it to descendants of $V_{j-1,1}$. Finally, there should be a $v'_{j+1,1}$ which is technically a sub-singular vector, i.e. it is singular in the quotient by the representation generated by $V_{j,1}$ (in other words, positive modes of Virasoro send it to descendants of $V_{j,1}$). This is schematically represented in Fig. 5, where arrows indicate action of the Virasoro algebra. Nodes connected by southeast arrows contribute $K_{1,1+2j}$ and $K_{1,1+2(j-1)}$ to the character; these correspond to a submodule and a quotient module, both of which are isomorphic to standard modules; the head and foot of each of these are indicated by the open and closed circles. In the CFT, these vectors that are highest weights modulo submodules will be the closest that we can come to finding primary fields for the Virasoro algebra.

Representations with the quartet structure such as $\mathcal{R}_j$ have been abstractly studied before and are called staggered modules in [30]. It can be shown that they depend on one characteristic parameter, which can be defined through the equation

$$\begin{aligned} L_1^{j-1} V'_{j,1} &= \beta_j v_{j-1,1}, \\ L_k V'_{j,1} &= 0, \quad k \geq 2. \end{aligned} \quad (3.18)$$

We do not know the relevant values of β_j in our models; it would be interesting to obtain them from the spin-1/2 chain, or from the symplectic fermion field theory discussed below. Presumably our modules are special as they are “projective.”

Now we interpret the continuum limit using the symplectic fermion representation of the theory. This follows from writing the spin-1/2 chain, or the supersymmetric formulation for $n = 1$, as a free-fermion system. The continuum limit can then be taken. As we consider the open chain case, one must treat the boundary conditions on the fermion fields. The fermion fields can be viewed as a complex scalar fermion, and its conjugate. These fermions are partners under a supersymmetry (graded Lie algebra) of free bosons, and the corresponding continuum free boson theory is a complex scalar boson. The appropriate boundary conditions for the latter were considered in detail in Ref. [42], both for the lattice model (discrete in both space and time directions, and called a network model) and in the

continuum. Hence the boundary condition on the fermions also is a variant of a Neumann boundary condition, in which on the boundary, the derivative of the field at some fixed angle (the “Hall angle” [42]) to the normal to the boundary must vanish (the Hall angle is a free parameter of the theory). This is due to the alternation of the two types of site, in the fundamental and its dual, along the chain when viewed as a representation of $\mathfrak{gl}(1|1)$. A property of these boundary conditions is that, as for ordinary Neumann boundary conditions, when we find the normal modes of the continuum theory on a finite interval, there is a zero mode for the fermion. Indeed, the entire spectrum is independent of the choice of Hall angle. For any value of the Hall angle, the theory can be mapped to the Hall angle zero case, which is the ordinary Neumann boundary condition. The transformation is a dual rotation, mapping the field to a linear combination of itself and its dual field; this transformation is nonlocal in terms of the fields, but linear, and commutes with the $U(1)$ symmetry of the complex fermion theory. After this transformation, the fields of the theory are the fermion fields ψ with two real components ψ_1, ψ_2 , where $\psi = \psi_1 + i\psi_2$, and their derivatives with respect to z only (as they are boundary fields), and products of these. The Lagrangian density is $\frac{1}{2}(\partial_\mu \psi_1 \partial_\mu \psi_2 - \partial_\mu \psi_2 \partial_\mu \psi_1)$, and thanks to the Neumann boundary condition, there is a well-known exact symplectic $[\mathrm{Sp}(2, \mathbf{R})]$ symmetry of this theory, that acts locally on the fields. Note however that for the original field variables that were natural for the spin chain, only the $U(1)$ subgroup of this symmetry acts locally (it corresponds to S^z or fermion number symmetry in the chain), while the other two generators of the $\mathfrak{sp}_2$ Lie algebra are nonlocal in those variables and arise as the “renormalized squares” of S^+ and S^- in the $U_q(\mathfrak{sl}_2)$ language, or as some of the nonlocal $k = 2$ basis elements constructed in Ref. [22].

Now we classify these local operators according to the Virasoro [and $U_q(\mathfrak{sl}_2)$] module structure we have worked out. The arrows connecting simple modules in the indecomposables of either algebra are always due to the fermion zero modes [43], which correspond to the local fields ψ_i without derivatives, as in operator products $\partial\psi_1(z)\psi_2(0) \sim 1/z$, while ψ_i without a derivative cannot be produced by any operator products of the local fields if it is not already a factor in an operator. The case $j = 2$ is typical. Then $h_{j,1} = 1$, the vector $V_{j,1}$ can be identified e.g. as the derivative of either symplectic fermion, say $V_{j,1} = \partial\psi_2$. The vector $v_{1,1} = \psi_2$ has dimension $h_{1,1} = 0$ and is a Virasoro highest weight: it generates a reducible representation, in which $\partial\psi_2$ appears as a singular vector. We still need a field $v'_{3,1}$ which is a combination involving $\psi_1\partial\psi_2\partial^2\psi_2$ etc. It is not primary, nor is $v'_{2,1} = \psi_1\psi_2\partial\psi_2$. By interchange $\psi_1 \leftrightarrow \psi_2$, two such Virasoro modules appear, which corresponds to the dimension of the $U_q(\mathfrak{sl}_2)$ simple module, which is just 2.

For the case $j = 1$, the cell lacks its left hand side and looks as on figure 6. One has for instance $L_{-1}(\psi_1\psi_2) = \partial(\psi_1\psi_2)$ while $L_1(\partial(\psi_1\psi_2)) = 1$, etc.

For the $U_q(\mathfrak{sl}_2)$ structure, we may notice that general local fields that are highest weights for Virasoro, modulo submodules, can be written as one of the following form: (i) $\mathcal{S}(\partial\psi_{i_1}\partial^2\psi_{i_2}\cdots\partial^{j-1}\psi_{i_{j-1}})$, with $j = 1, 2, \dots$, where the symmetrizer $\mathcal{S}$ symmetrizes in the indices $i = 1, 2$; (ii) the same with ψ_{i_0} inserted inside the symmetrizer $\mathcal{S}$; (iii) a similar expressions as in (ii) but with the index on ψ_{i_0} contracted with one of the $\partial^k\psi_i$ using the symplectic form; (iv) the form (i) times $\psi_1\psi_2$. For each j , these all have the same scaling dimension (meaning the eigenvalue of L_0 , modulo submodules). Those of form (i) are the foot of the Virasoro module $\mathcal{R}_j$, that is they form a simple module of scaling dimension $j(j-1)/2$; this may be seen using operator products with the stress tensor $T = -\frac{1}{2}(\partial\psi_1\partial\psi_2 - \partial\psi_2\partial\psi_1)$. There are j distinct such operators, because of the symmetrization, and so they form a simple $U_q(\mathfrak{sl}_2)$ module, with the $U_q(\mathfrak{sl}_2)$ spin $j-1$. The local fields of form (ii) in which ψ_{i_0} are included form a multiplet of dimension $j+1$, those of type (iii) form a multiplet of dimension $j-1$, and finally those of form (iv) in which $\psi_1\psi_2$ is included form a multiplet of dimension j . For each j , these four types of local operators (or three types if $j = 1$) make up the tilting module T_j of $U_q(\mathfrak{sl}_2)$, listed here starting with the foot and ending with the head. The symplectic or $\mathfrak{sp}_2$ symmetry of symplectic fermions acts to map the fields of the same type into each other; if we view the boundary symplectic fermion theory as a chiral theory in the plane, then there are three corresponding Noether currents $\mathcal{S}(\psi_i\partial\psi_{i'})$. The doublet of currents $\partial\psi_i$ of $\mathfrak{psl}(1|1)$ [20] act (through operator product) to remove an undifferentiated ψ_i from the boundary local fields. These are the arrows in the $U_q(\mathfrak{sl}_2)$ quartets. The $U_q(\mathfrak{sl}_2)$ algebra can here be viewed as isomorphic to a Lie superalgebra generated by the line integrals of these five local currents (which also correspond to the five generating elements of $U_q(\mathfrak{sl}_2)$ at a root of unity); this Lie superalgebra is not semisimple (as a Lie superalgebra), because the $\mathfrak{psl}(1|1)$ generators form an ideal.




Figure 6: Indecomposable representation for $j = 1$.

For the Virasoro structure, it suffices to notice that the stress tensor T also maps each of the four types into themselves by operator products, and also can reduce the number of undifferentiated fermion fields in a local operator by one or two. These actions then exhibit the commuting actions of Virasoro and $U_q(\mathfrak{sl}_2)$, and show how the staircase diagram structure appears in terms of the local fields.

The partition function can also be interpreted in terms of symplectic fermions. In terms of the characters of simple Virasoro modules and dimensions of simple modules of $U_q(\mathfrak{sl}_2)$, it becomes

$$Z = \sum_{j=0}^{\infty} j(2\chi_{j,1} + \chi_{j-1,1} + \chi_{j+1,1}) = \frac{2\theta_2(\hat{q})}{\eta(\hat{q})} = \det(-\mathcal{D}_{A,N}), \quad (3.19)$$

where $\mathcal{D}_{A,N}$ is the Laplacian on the strip with Neumann boundary conditions in the space direction and antiperiodic boundary conditions in the time direction. For symplectic fermions, the vanishing of the “unmodified” (or quantum trace) partition function can be viewed as due to the fermion zero modes, as is well known (in the present language, it is the determinant of the Laplacian with periodic boundary conditions in the time direction, and this Laplacian has a zero mode).

We now discuss fusion of the tilting modules T_j (of dimension $4j$) labelled by j , in other words the Clebsch-Gordan series for $U_q(\mathfrak{sl}_2)$ at $q = i$. A tensor product of indecomposable tilting modules necessarily decomposes into a direct sum of tilting modules, and given this fact, the decomposition can be easily determined by using characters [40]. It takes the form:

$$T_{j_1} \otimes T_{j_2} = T_{j_1+j_2} \oplus 2T_{j_1+j_2-1} \oplus \dots \oplus 2T_{|j_1-j_2|+1} \oplus T_{|j_1-j_2|}. \quad (3.20)$$

Here $2T$ means $T \oplus T$, and for $j_1 = j_2$, the term T_0 is absent; for the present we may view this exception as handled by our convention that $T_j = 0$ if $j < 1$. (A check on the result can be easily obtained by considering the dimensions $2j+1 + (2j-1)$ of the tilting modules.) It follows from Morita equivalence that the same fusion rules apply to the tilting modules of our supersymmetry algebras $\mathcal{A}_{n|n}$ also [22].

From the general theory [22], the fusion rules for the product *functor* for the indecomposable *projective* TL modules have the same form,

$$P_{j_1} \times P_{j_2} = P_{j_1+j_2} + 2 \sum_{j=|j_1-j_2|+1}^{j_1+j_2-1} P_j + P_{|j_1-j_2|}, \quad (3.21)$$

and so do those for the projective Virasoro modules in the continuum limit:

$$\mathcal{R}_{j_1} \times \mathcal{R}_{j_2} = \mathcal{R}_{j_1+j_2} + 2 \sum_{j=|j_1-j_2|+1}^{j_1+j_2-1} \mathcal{R}_j + \mathcal{R}_{|j_1-j_2|}. \quad (3.22)$$

(Again, recall that the modules P_0 and $\mathcal{R}_0$ are zero.) The summations may be re-organized into the form

$$\mathcal{R}_{j_1} \times \mathcal{R}_{j_2} = \sum_j' (2\mathcal{R}_j + \mathcal{R}_{j-1} + \mathcal{R}_{j+1}), \quad (3.23)$$

where the sum is over $j = j_1 + j_2 - 1, j_1 + j_2 - 3, \dots, |j_1 - j_2| + 1$. This shows that these results agree with those for $c = -2$ in Ref. [44]. There, some indecomposables of Virasoro are built somewhat explicitly, and fusion is defined and calculated up to level six through the algorithm in [45]; the final results are then conjectured. The correspondence in the latter reference would be $m \equiv j$ and $\mathcal{R}_{m,1} \equiv \mathcal{R}_j$. Calculations in [44] are extremely technical, and it is quite remarkable that the results can be derived from fusion of indecomposables of $U_q(\mathfrak{sl}_2)$.

We have concentrated here on the fusion rules for tilting modules on the $U_q(\mathfrak{sl}_2)$ side, and the projective modules on the TL or Virasoro side, as these are the most natural starting point, and there is an equivalence between these fusion products. One can also derive from these the fusion rules for some other types of modules, such as the standard and simple modules. One must be careful in using the $U_q(\mathfrak{sl}_2)$ fusion results on the TL/Virasoro side as the equivalence may not always hold for the non-tilting modules. The general definition of fusion for the TL or Virasoro modules is always the module induced from the tensor product by the inclusion of the algebras for two chains into that for a single chain, by joining them end to end.

We emphasize that these fusion rules can be viewed as describing operator products of the conformal fields that transform as Virasoro modules discussed here, for example for the symplectic fermions, for which the “pseudoprimary” fields were written down earlier, each of which is the primary field in a simple subquotient module. They also apply to the corresponding fields in the supersymmetric models, where the fields have different multiplicities given by the dimensions D_j^0 . In applications the fusion rules given here will have to be combined with those for the symmetry algebra $[U_q(\mathfrak{sl}_2) \text{ or } \mathcal{A}_{n|n}]$, as the fields transform under both algebras. These combined forms can be obtained in the finite- L lattice models, where they come from decomposing tensor products simply as vector spaces (the multiplicities from the symmetry algebra side make the dimensions of spaces conserved), and then the continuum limit can be taken.

As a final note, we mention that in the case of unoriented loops, we should consider odd as well as even values of $2L$. Then the half integer spins j , for which $2j$ is odd also appear in the spectrum. For $q = i$, these standard modules remain simple, both for $U_q(\mathfrak{sl}_2)$ and (hence) also for TL. They appear then as extra circles interspersed among the integer j values in the staircase diagrams, with no extra explicit arrows, so these modules are both tilting and projective also. Fusion rules for products involving these can be found along the same lines as for j_1, j_2 integer (the fusion rules for a product of j_1 with j_2 both integers are unchanged, and still close on integer j only). In terms of dense polymers, the unoriented-loops model contains local boundary fields for odd as well as even numbers of legs, the number of legs being (as always for the loop models) $2j$ for fields in a standard module of spin j .

4 The case $m = 1$ ($c = 0$) — percolation

In this section we follow a similar procedure as in the previous one, for the XXZ chain of even length $2L$ with $m = 1$, so $q = e^{i\pi/3}$. In this case the loop model describes dense, oriented loops which can be identified with the boundaries of clusters in percolation at the critical percolation threshold. In the spin chains with $\mathfrak{gl}(n+1|n)$ supersymmetry [20], the continuum limit in this case corresponds to the fixed-point theory of the $\mathbf{CP}^{n|n}$ nonlinear sigma model at $\theta = \pi$, with the enlarged symmetry algebra $\mathcal{A}_{n+1|n}$ that is Morita equivalent to $U_q(\mathfrak{sl}_2)$ with $q = e^{i\pi/3}$ [22]. The $n = 1$ case is thus connected with the spin quantum Hall transition also [3]. Because the TL algebra for $-m$ is isomorphic to that for m , the first, algebraic part of the analysis is the same for the $m = -1$ finite- L spin chain for that case also, up to a change in sign of the Hamiltonian. Because of the latter sign change, a distinct continuum limit is obtained for $m = -1$ which is not considered in this section. We note that since the chain has even length, only integer spins j occur, and for these the symmetry algebras $U_q(\mathfrak{sl}_2)$ and $U_{-q}(\mathfrak{sl}_2)$ are isomorphic.

We start with the analysis of the spin-1/2 chain under $U_q(\mathfrak{sl}_2)$. For $q = e^{i\pi/3}$, so $r = 3$, and the representations depend on $2j + 1$ modulo 3, instead of modulo 2 as in the previous case. The standard modules for $j \geq 0$ with $j \equiv 1 \pmod{3}$ remain irreducible. The remaining standard modules for $j > 0$ become reducible in the manner described for all roots of unity, and the corresponding tilting modules (which are the direct summands in the spin-1/2 chain) contain a submodule and a quotient module

that are standard, and which have adjacent j -values if the values $j \equiv 1 \pmod{3}$ are skipped. Similarly, the j values assigned to the simple subquotient modules of the standard modules are also adjacent if the values $j \equiv 1 \pmod{3}$ are skipped, and the value at the head is the same as the value for the standard module. The standard module for $j = 0$ is one dimensional, and so is simple, and also appears as a summand in the chain. Thus the tilting modules have the following structure [40, 25, 26]:

$$T_j : \left\{ \begin{array}{ll} \begin{array}{c} j-1 \\ \swarrow \quad \searrow \\ j-3 \quad j \\ \searrow \quad \swarrow \\ j-1 \end{array} & j \equiv 0 \pmod{3}, \\ j & j \equiv 1 \pmod{3}, \\ \begin{array}{c} j-2 \\ \swarrow \quad \searrow \\ j-3 \quad j \\ \searrow \quad \swarrow \\ j-2 \end{array} & j \equiv 2 \pmod{3}, \end{array} \right.$$

for $j = 0, 1, \dots, L$. Here again, simple subquotient modules labeled $j' \leq 0$ are understood to be zero and can be omitted. In the reducible modules, the pairs connected by southwest arrows form standard modules. Given this structure and the dimensions $2j + 1$ for the standard modules, the dimensions of the simple modules can be found recursively. Introducing $p = \lfloor j/3 \rfloor$ (where $\lfloor x \rfloor$ is the largest integer $\leq x$), the dimensions of the simple modules are $2p+1, 6p+1, 4p+4$ for $j \equiv 0, 1, 2, \pmod{3}$ respectively ($p \geq 0$). Thus in terms of dimensions of the simple subquotients, we have for the two non-trivial classes:

$$T_{3p} : \begin{array}{c} (4p) \\ \swarrow \quad \searrow \\ (2p-1) \quad (2p+1) \\ \searrow \quad \swarrow \\ (4p) \end{array} \quad T_{3p+2} : \begin{array}{c} (2p+1) \\ \swarrow \quad \searrow \\ (4p) \quad (4p+4) \\ \searrow \quad \swarrow \\ (2p+1) \end{array}$$

The q -dimensions, and the dimensions and superdimensions of the standard and tilting modules over $\mathcal{A}_{n+1|n}$, can be calculated similarly, as we showed in the case of $m = 0$. In particular, the q -dimensions (superdimensions) D_j^+ of all the tilting modules over $U_q(\mathfrak{sl}_2)$ (resp., $\mathcal{A}_{n+1|n}$) are zero except for the $j = 0$ tilting module, which has $D_0^+ = 1$.

Taking $L = 3$ as an example, the decomposition of the Hilbert space of the chain under $U_q(\mathfrak{sl}_2)$ is, in terms of spins j ,

$$0 \times 1, \quad 1 \times 9, \quad \begin{array}{c} 0 \\ \searrow \\ 2 \\ \swarrow \\ 0 \end{array} \times 4, \quad \begin{array}{c} 2 \\ \swarrow \quad \searrow \\ 0 \quad 3 \\ \searrow \quad \swarrow \\ 2 \end{array} \times 1,$$

for a total of 64 states indeed. The staircase diagram is shown in Fig. 7.

For the TL algebra, the direct summands, with their multiplicities, are [24, 25, 26]

$$\begin{array}{c} (4) \\ \swarrow \quad \searrow \\ (1)' \quad (1) \\ \searrow \quad \swarrow \\ (4) \end{array}, \quad (9) \times 3, \quad \begin{array}{c} (1) \\ \swarrow \\ (4) \\ \searrow \\ (1) \end{array} \times 4, \quad (1) \times 3, \quad (4.1)$$

in order of increasing j . The dimensions of the simple subquotients are the multiplicities of the $U_q(\mathfrak{sl}_2)$ tilting modules, and the multiplicities are the dimensions of the $U_q(\mathfrak{sl}_2)$ simple modules. But notice




Figure 7: The structure of the spin-1/2 chain for $q = e^{i\pi/3}$ and $2L = 6$ sites, as a representation of $U_q(\mathfrak{sl}_2) \otimes \text{TL}_{2L}(q)$, or the same for the $\mathfrak{gl}(1+n|n)$ -supersymmetric chain with $2L = 6$ sites as a representation of $\mathcal{A}_{1+n|n}(2L) \otimes \text{TL}_{2L}(q)$.

that dimension 1 appears in connection with two different summands under $U_q(\mathfrak{sl}_2)$, and these are not expected to be isomorphic as TL modules—they have different j values. For this reason, one of them has been distinguished here by writing $(1)'$. All of these indecomposable modules are projective, except for the TL singlet at the end. However, there must also be an additional projective module, which has the simple module $(1)'$ at the top, and must “cover” the lower part of the TL tilting module in which it occurs. This remaining indecomposable projective module thus has the form

$$\begin{array}{c} (1)' \\ \searrow \\ (4) \end{array} \quad (4.2)$$

(and is a standard module), for a total of $L + 1 = 4$ isomorphism classes as expected here, since they must correspond to the four distinct indecomposable tilting modules of $U_q(\mathfrak{sl}_2)^{(4)}$.

For general L , the pattern is similar. There is always a singlet standard module with spin $j = L$ which is a TL tilting module. For $L \not\equiv 1 \pmod{3}$, it is not projective. For $L \equiv 1 \pmod{3}$, this singlet is projective as well as tilting. But then there is always a simple summand (tilting module) at the largest $j \not\equiv 1 \pmod{3}$, which is not a TL projective module. The remaining indecomposable tilting modules [including those standard modules with $j \equiv 1 \pmod{3}$ which are simple] are also projective, and there is one additional projective module for $j = 0$, that has a (different) singlet at the top. Thus




Figure 8: Space of states in the continuum $m = 1$ theory, showing commuting actions of Virasoro and $U_q(\mathfrak{sl}_2)$ (or $\mathcal{A}_{1+n|n}$).

the projective modules for TL are [24, 25, 26], in terms of spin labels,

$$P_j : \left\{ \begin{array}{ll} \begin{array}{c} 0 \\ \searrow \\ 2 \end{array} & j = 0, \\ \begin{array}{c} j \\ \swarrow \quad \searrow \\ j-1 \quad j+2 \\ \swarrow \quad \searrow \\ j \end{array} & j \equiv 0 \pmod{3} \text{ and } j > 0, \\ j & j \equiv 1 \pmod{3}, \\ \begin{array}{c} j \\ \swarrow \quad \searrow \\ j-2 \quad j+1 \\ \swarrow \quad \searrow \\ j \end{array} & j \equiv 2 \pmod{3}, \end{array} \right.$$

where again simple modules with $j < 0$ or $j > L$ are understood to be zero. The standard sub- and quotient modules are the pairs connected by a southeast arrow.

Once this structure is known, the dimensions d_j^0 of the simple modules can be found recursively. For $j \equiv 0$, the standard module is simple, and has dimension $d_j^0 = d_j$ as before. For the other cases, one has $d_j^0 = d_j - d_{j+2}^0$ [$j \equiv 0 \pmod{3}$] and $d_j^0 = d_j - d_{j+1}^0$ [$j \equiv 2 \pmod{3}$]. These can be solved starting from $d_L = d_L^0 = 1$. It is possible to show that $d_0^0 = 1$ for all L (as already stated above), and then the recursion can also be solved starting from the $j = 0$ end. For $q = e^{i\pi/3}$, we have not found any very illuminating closed-form expressions for the dimensions d_j^0 for $j > 0$.

In the large L limit, the staircase diagram continues to infinity, and resembles that for $q = i$, except that extra simple modules appear at $j \equiv 1 \pmod{3}$, and there is now a node at $(0, 0)$; see Fig. 8.

Let us now consider the continuum limit in more detail. For $c = 0$, the Kac formula reads

$$h_{rs} = \frac{(3r - 2s)^2 - 1}{24}. \quad (4.3)$$

The conformal weights appearing of the primary fields appearing as subquotients in the lattice partition function are

$$h_{1,1+2j} = \frac{j(2j-1)}{3}. \quad (4.4)$$

We again consider only the case j integer, and often abbreviate $h_{1,1+2j} = h_j$. For $j \not\equiv 1 \pmod{3}$, these weights are integers.

Introducing the same notation $K_{1,1+2j} \equiv K_j$ for the Virasoro characters of the standard modules, which we again denote $r_{1,1+2j} \equiv r_j$, we have

$$K_j \equiv K_{1,1+2j} = \frac{\hat{q}^{h_{1,1+2j}} - \hat{q}^{h_{1,-1-2j}}}{P(\hat{q})}, \quad (4.5)$$

and from the limit of the spin chain we expect to have the relations (using again $p = \lfloor j/3 \rfloor$)

$$\begin{aligned} K_{3p} &= \chi_{3p} + \chi_{3p+2}, \\ K_{3p+1} &= \chi_{3p+1}, \\ K_{3p+2} &= \chi_{3p+2} + \chi_{3p+3}, \end{aligned} \quad (4.6)$$

where χ_j are the characters of the associated irreducible Virasoro representation with highest weight $h_{1,1+2j}$ (denoted here by R_j). As for $m = 0$, this expectation is correct as it agrees with the known structure of Virasoro representations in the Kac table for $c = 0$ [33, 46]; the standard modules r_j contain a unique simple submodule when $j \not\equiv 1 \pmod{3}$. The irreducible characters are known, and take different forms depending on $j \pmod{3}$ [46]. If $j = 3p + 1$, we have

$$\chi_{j=3p+1} = \hat{q}^{(3+12p)^2/24} \frac{1 - \hat{q}^{3+6p}}{P(\hat{q})}. \quad (4.7)$$

If $j = 3p + 2$, we have

$$\chi_{j=3p+2} = \sum_{n \notin [-2p-1, -1]} \frac{\hat{q}^{(12n+12p+7)^2/24} - \hat{q}^{(12n+12p+11)^2/24}}{P(\hat{q})}, \quad (4.8)$$

and if $j = 3p$

$$\chi_{j=3p} = \sum_{n \notin [-2p, -1]} \frac{\hat{q}^{(12n+12p-1)^2/24} - \hat{q}^{(12n+12p+7)^2/24}}{P(\hat{q})}. \quad (4.9)$$

In particular, the irreducible character of the identity is $\chi_0 = 1$.

For $j = 3p + 1$, the standard representation $r_{j=3p+1}$ is irreducible, and we can view it as our projective module. For $j \not\equiv 1 \pmod{3}$ we expect that there are indecomposable projective modules $\mathcal{R}_j$

that are the limits of the TL modules, and correspond to them in submodule structure, thus

$$\mathcal{R}_j : \left\{ \begin{array}{ll} \begin{array}{c} R_0 \searrow \\ R_2 \end{array} & j = 0, \\ \begin{array}{c} R_j \searrow \\ R_{j-1} \searrow \quad R_{j+2} \searrow \\ \mathcal{R}_j \end{array} & j \equiv 0 \pmod{3} \text{ and } j > 0, \\ R_j & j \equiv 1 \pmod{3}, \\ \begin{array}{c} R_j \searrow \\ R_{j-2} \searrow \quad R_{j+1} \searrow \\ R_j \end{array} & j \equiv 2 \pmod{3}, \end{array} \right.$$

for $j = 0, 1, \dots$. Apart from the the $j = 0$ case, and the intruding $j \equiv 1 \pmod{3}$ simple modules, these resemble the projective modules of the $c = -2$ case. The characters of the non-trivial (non-standard) cases are clearly, for $j = 3p$,

$$\text{Tr}_{\mathcal{R}_j} \hat{q}^{L_0 - c/24} = K_{3p} + K_{3p-1} = 2\chi_{3p} + \chi_{3p-1} + \chi_{3p+2}, \quad (4.10)$$

and for $j = 3p + 2$, similarly,

$$\text{Tr}_{\mathcal{R}_j} \hat{q}^{L_0 - c/24} = K_{3p+2} + K_{3p} = 2\chi_{3p+2} + \chi_{3p} + \chi_{3p+3}. \quad (4.11)$$

The partition function of the continuum limit of the spin-1/2 chain is again given by

$$Z = \sum_{j=0}^{\infty} (2j+1) K_j = \sum_{j=0}^{\infty} (2j+1) \frac{\hat{q}^{h_{1,1+2j}} - \hat{q}^{h_{1,-1-2j}}}{P(\hat{q})}. \quad (4.12)$$

For the q -trace instead of the trace, one replaces $2j+1$ by $[2j+1]_q$, and the resulting formula also gives the “unmodified” partition function of the supersymmetric models with symmetry algebra $\mathcal{A}_{n+1|n}$ [20]. For the partition function of the latter, one instead replaces $2j+1$ by the dimensions $D'_j = [2j+1]_{q'}$, where $q' + q'^{-1} = 2n+1$ for these $m = 1$ models (see Appendix B in ref. [20]).

Understanding the representations $\mathcal{R}_j$ more explicitly is more difficult than for $m = 0$, since we do not have an explicitly solvable version analogous to the symplectic fermions. One might think of using the Feigin-Fuchs free field representation of the theory properly extended by some extra space to allow for the presence of Jordan cells, along the lines of [14]. We have not done so however, but suspect that the continuum limits of the present models are different anyhow, since they involve a single copy of the $h = 0$ field. This will be discussed in more detail when we turn to the case of the torus in a later paper.

We can now study fusion of the $U_q(\mathfrak{sl}_2)$ tilting modules T_j . The decomposition of a tensor product of T_{j_1} with T_{j_2} has to be considered in six cases, as it depends on j_1 and $j_2 \pmod{3}$. We let $p_1 = \lfloor j_1 \rfloor / 3$, $p_2 = \lfloor j_2 \rfloor / 3$. For cases in which $j_1, j_2 \not\equiv 1 \pmod{3}$, we have:

$$T_{3p_1+2} \otimes T_{3p_2+2} = \bigoplus_r' \left(2T_{3r+2} \oplus \underline{T_{3r}} \oplus T_{3r+3} \oplus T_{3r+1} \oplus \underline{T_{3r+1}} \oplus \underline{T_{3r-2}} \oplus T_{3r+4} \right), \quad (4.13)$$

where the sum runs over $r = |p_1 - p_2|, |p_1 - p_2| + 1, \dots, p_1 + p_2$. Here and in the following, underlined terms are absent if $r = 0$. Similarly

$$T_{3p_1} \otimes T_{3p_2} = \bigoplus_r' \left(2T_{3r+2} \oplus \underline{T_{3r}} \oplus T_{3r+3} \oplus 4T_{3r+1} \right), \quad (4.14)$$

where the sum runs over $r = |p_1 - p_2|, |p_1 - p_2| + 1, \dots, p_1 + p_2 - 1$. Finally, for the mixed fusion

$$T_{3p_1+2} \otimes T_{3p_2} = \bigoplus_r' (2T_{3r} \oplus T_{3r-1} \oplus T_{3r+2} \oplus 2T_{3r-2} \oplus 2T_{3r+1}), \quad (4.15)$$

where the sum runs over $r = p_1 - p_2 + 1, p_1 - p_2 + 2, \dots, p_1 + p_2$ if $p_1 \geq p_2$ and $r = p_2 - p_1, p_2 - p_1 + 1, \dots, p_1 + p_2$ if $p_1 < p_2$.

For the cases in which (without loss of generality) $j_1 \equiv 1 \pmod{3}$, we have

$$T_{3p_1+1} \otimes T_{3p_2+1} = \bigoplus_r' (T_{3r+1} \oplus T_{3r+2}), \quad (4.16)$$

where the sum runs over $r = |p_1 - p_2|, |p_1 - p_2| + 1, \dots, p_1 + p_2$. Moreover

$$T_{3p_1+1} \otimes T_{3p_2+2} = \bigoplus_r' (2T_{3r+1} \oplus \underline{T_{3r}} \oplus T_{3r+3}), \quad (4.17)$$

where the sum runs over $r = |p_1 - p_2|, |p_1 - p_2| + 1, \dots, p_1 + p_2$, and

$$T_{3p_1+1} \otimes T_{3p_2} = \bigoplus_r' (2T_{3r-1} \oplus \underline{T_{3r-2}} \oplus T_{3r+1}), \quad (4.18)$$

where the sum runs over $r = p_1 - p_2 + 1, p_1 - p_2 + 2, \dots, p_1 + p_2$ if $p_1 \geq p_2$ and $r = p_2 - p_1, p_2 - p_1 + 1, \dots, p_1 + p_2$ if $p_1 < p_2$.

Finally, we note the special case $j_1 = 0$, when

$$T_0 \otimes T_j = T_j \quad (4.19)$$

for all $j \geq 0$.

The fusion rules for the Virasoro projective modules follow immediately from the correspondence $T_j \leftrightarrow \mathcal{R}_j$ (and also those for TL projective modules, from the correspondence $T_j \leftrightarrow P_j$). As far as we know, these fusion rules for a logarithmic theory at $c = 0$ are new. This is one of the central results of this paper.

We note that the $j = 0$ projective module $\mathcal{R}_0$ does indeed behave as the identity in the fusion rules, as expected from the relation with $U_q(\mathfrak{sl}_2)$, because it contains the identity and the stress tensor, which map any of the projective modules $\mathcal{R}_j$ for $j > 0$ into themselves. None of the fusion rules produce the module T_0 (on the symmetry side) or $\mathcal{R}_0$ (on the Virasoro side). The identity field, and the stress tensor, also occur within the T_2 (resp., $\mathcal{R}_2$) modules, so that conformally-invariant correlation functions do exist.

5 The case $m = 0$ for dilute loops — critical polymers

A lattice (vertex) model for dilute loops can be formulated on the square lattice with a two-colour TL algebra [47], or a spin-1 chain with the states $S_z = \pm 1$ viewed as occupied by parts of loops, and $S_z = 0$ as empty. It does have $U_q(\mathfrak{sl}_2)$ symmetry, with the strands transforming as spin-1/2. From a symmetry point of view, the models are identical to the dense cases. This is also true within the supersymmetric formulation [20], in which the obvious supersymmetry algebra is (the enveloping algebra of) $\mathfrak{osp}(m + 2n|2n)$, and acts on the occupied sites as the vector (defining) representation, and trivially on the empty sites. As the models forbid the strands to cross, the global symmetry algebra is actually larger than $\mathfrak{osp}(m + 2n|2n)$ [22]. We will consider here the unoriented version of dilute loops with $m = 0$, for variety and because this is of natural interest for polymers. This means the number of strands on any time-slice of the system is unrestricted. Because of the existence of empty sites, both parities of $2j$ occur for each length of system. For $m = 0$, this model possesses both a high-temperature phase and a low-temperature phase, separated by a transition. The continuum limit in the low-temperature phase is in the same universality class as the (unoriented loops) $m = 0$ dense polymer theory above. The critical point describes dilute polymers. It is the same universality




Figure 9: Space of states in the continuum dilute $m = 0$ theory, showing commuting actions of Virasoro and $U_q(\mathfrak{sl}_2)$ (or $\mathcal{B}_{n|n}$).

class as in an $\mathfrak{osp}(2n|2n)$ -invariant ϕ^4 (Landau-Ginzburg) formulation of dilute polymers, or as in an $\mathfrak{osp}(2n|2n)$ -invariant nonlinear sigma model with target space a supersphere, $S^{2n-1|2n}$ [20].

We will not define in detail here the lattice model, but we make some comments. For all q , the algebra generated by the nearest-neighbor interactions in the spin-1 chain that represents dilute loops is clearly somehow similar to the TL algebra, even though the dimensions of the algebras are different. We suspect that this relation is in fact Morita equivalence (with a partial exception that will be mentioned below): the algebra generated by the spin-1 chain is Morita equivalent to the TL algebra (generated by the nearest-neighbor terms in the spin-1/2 chain) for the same number of sites, and at the same q . This is motivated by the common symmetry structure for all q , and by the relation of this with the TL (or Virasoro) structure. It remains a conjecture as we have not established that the spin-1 model and $U_q(\mathfrak{sl}_2)$ form a “dual pair”. The exception occurs when $m = 0$, since as we will see the singlet module at $(0, 0)$ is present in the spin-1 case, but was absent for the spin-1/2 model. This is connected with a well-known phenomenon in these loop models: The dilute model possesses a low- T phase that is in the same universality class as the dense model. Yet the dilute model has non-vanishing partition function (here it is the “unmodified” partition function, defined using the q -trace in the transfer matrix approach, that we mean), but that of the dense model vanishes for $m = 0$. The discrepancy is resolved because the state with no polymer loops exists in the dilute model, and produces the nonzero partition function, but there is no such configuration in the dense model. In the dense phase of the dilute model, this state occurs at high energy and is dropped in the continuum limit, so the partition function (in which energies of the spin chain are measured from the ground state energy) becomes zero. This configuration with no loops corresponds to the state of the chain at $(0, 0)$ in the staircase diagram for $m = 0$.

As we do not consider the lattice algebraic formulation in detail here, we pass directly to the continuum theory at the critical point for $m = 0$. The staircase diagram is shown in Fig. 9. It has the same form as the dense polymer case, except that the circle at $(0, 0)$ is now included. The $U_q(\mathfrak{sl}_2)$ structure is the same as for dense polymers, and we include the $2j$ odd cases for the unoriented case. Here the singlet tilting module T_0 does appear with nonzero multiplicity in the chain. For the supersymmetric models, the symmetry algebra for the unoriented loops case is called $\mathcal{B}_{n|n}$ in Ref. [22].

On the Virasoro side, the conformal weights are taken from the first *row* of the $c = 0$ Kac table, so

$$h_j \equiv h_{1+2j,1} = \frac{j(3j+1)}{2}. \quad (5.1)$$

The characters of the standard modules read

$$K_j = \frac{\hat{q}^{h_{1+2j,1}} - \hat{q}^{h_{-1-2j,1}}}{P(\hat{q})}. \quad (5.2)$$

Using standard character formulas we check the identities

$$\begin{aligned} K_{p+1/2} &= \chi_{p+1/2}, \\ K_p &= \chi_p + \chi_{p+1}, \end{aligned} \quad (5.3)$$

similar to the $m = 0$ dense case. The partition function of the continuum limit of the spin one chain is again given by

$$Z = \sum_{j=0}^{\infty} (2j+1) \frac{\hat{q}^{h_{1+2j,1}} - \hat{q}^{h_{-1-2j,1}}}{P(\hat{q})}. \quad (5.4)$$

For the q -trace instead of the trace, one replaces $2j+1$ by $D_j = [2j+1]_q$ with $q = i$ (thus $D_j = (-1)^j$ for j integer, $D_j = 0$ for j half-integer), which also gives the unmodified partition function of the supersymmetric models with symmetry algebra $\mathcal{B}_{n|n}$ [22]. The latter is simply $Z_{\text{unmod}} = 1$ by the Euler identity. For the “modified” partition function of the supersymmetric models, one instead replaces $2j+1$ by the dimensions $D'_j = [2j+1]_{q'}$, where $q' + q'^{-1} = 2n$ for these models (see Appendix B in ref. [20]). For $n = 1$, the supersymmetry algebra is isomorphic to $U_q(\mathfrak{sl}_2)$, as in the dense case.

Projective modules for Virasoro are also similar to the dense case in structure, but now there is also $\mathcal{R}_0$ for $j = 0$, which is just a standard module, similarly to the $m = 1$ (percolation) case. It has the identity operator at its head, and the stress tensor T as the highest weight in the foot submodule. These fields are also part of the $\mathcal{R}_1$ module, similarly to percolation ($m = 1$).

Fusion rules for $j > 0$ are as for dense polymers. In the fusion of tilting modules of $U_q(\mathfrak{sl}_2)$ (or projective modules of Virasoro) for j_1 with $j_2 = j_1$ (and $j_1 > 0$), the term T_0 (resp., $\mathcal{R}_0$) is again absent, even though the corresponding modules are nonzero; this applies for $2j_1$ odd as well as even. T_0 behaves as the identity for the fusion rules:

$$T_0 \otimes T_j = T_j \quad (5.5)$$

for all $j \geq 0$, and similarly for fusion of $\mathcal{R}_0$ with $\mathcal{R}_j$. We note that the identity field occurs within the $\mathcal{R}_1$ module (unlike the dense case), and so conformally-covariant correlation functions can be produced.

6 Physical discussion

In this section, some physical discussion is given, by writing the algebraic statements in terms of operator products. The discussion is mainly for $c = 0$, and makes contact with that of Gurarie and Ludwig [17, 18]. We also make some remarks on the role of fields of weight 1 in the boundary theories.

The case $c = 0$ has probably been the most discussed in the literature on logarithmic CFTs. There, the existence of the “ c -catastrophe” (i.e., poles in $1/c$ in OPEs as $c \rightarrow 0$) led Gurarie [16, 17, 18] to the introduction of a *logarithmic partner* t for the stress energy tensor, satisfying the OPEs (up to non-divergent terms):

$$\begin{aligned} T(z)T(0) &= \frac{2T(0)}{z^2} + \frac{T'(0)}{z} + \dots, \\ T(z)t(0) &= \frac{b}{z^4} + \frac{2t(0) + T(0)}{z^2} + \frac{t'(0)}{z} + \dots, \end{aligned} \quad (6.1)$$

Notice that T is primary, but has a vanishing two-point function, while t is not primary. The tt OPE itself is more complicated and starts with a logarithm:

$$t(z)t(0) = -\frac{2b \ln z}{z^4} + \dots \quad (6.2)$$

It follows from these equations that T, t form a size two Jordan cell for L_0 :

$$\begin{aligned} L_0 T &= 2T, \\ L_0 t &= 2t + T. \end{aligned} \quad (6.3)$$

Gurarie went on to suggest that T, t were part of a ‘‘SUSY’’ multiplet containing at least two weight-two primary operators with non-vanishing two-point functions [16, 17, 18].

The foregoing algebraic analysis of our $c = 0$ lattice models gives rise immediately to the existence of t . Indeed, for the continuum limit of the chain, the Virasoro tilting module that contains the identity and the stress tensor is

$$\mathcal{R}_2 : \quad \begin{array}{ccc} & R_2 & \\ \swarrow & & \searrow \\ R_0 & & R_3 \\ \searrow & & \swarrow \\ & R_2 & \end{array} \quad (6.4)$$

Each simple subquotient module R_j that appears is a highest weight module, so (by the correspondence of states and operators) there is a corresponding pseudoprimary field. All these states have multiplicity 1. Thus the pseudoprimaries consist of the identity with conformal weight $h_0 = 0$, two fields with weight $h_2 = 2$, and one with dimension $h_3 = 5$. One of those with dimension 2 (the bottom field, as it is primary) is naturally identified with the stress energy tensor T , and should form a Jordan cell with its logarithmic partner - here the top field (with conformal weight 2) which we identify with t . The whole indecomposable representation is generated by acting on t with the Virasoro algebra. We now see that there is a fourth member of the multiplet of fields, which we call τ , and which has conformal weight $h_3 = 5$. We recall that this module can be analyzed into a submodule that is standard, and which is generated by the descendants of 1, and a quotient module that is also standard, and the quotient module is generated by t .

The arrow from R_2 to R_0 corresponds to the fact that t is not primary, and can be mapped to the identity under the action of Virasoro generators - this is directly related with the existence of the number b in the second OPE (6.1). The arrow from R_2 to R_3 indicates that the second OPE in (6.1) must be extended to include the field of dimension 5:

$$T(z)t(0) \sim \frac{b}{z^4} + \frac{2t(0) + T(0)}{z^2} + \frac{t'(0)}{z} + \dots + z c_{Tt}^T \tau(0) \dots \quad (6.5)$$

where c_{Tt}^T is some coefficient. τ is not primary, but only maps to descendants of T under action of Virasoro with positive modes:

$$T(z)\tau(0) \sim \frac{c_{T\tau}^T T(0)}{z^5} + \dots \quad (6.6)$$

Modulo the Virasoro submodule generated by T in this way, the field τ is simply the null Virasoro descendent of t that occurs in the Virasoro standard module when $c = 0$; it must not be set to zero.

The possible supersymmetry multiplet of fields of weight 2 [16, 17, 18] may be compared with the $U_q(\mathfrak{sl}_2)$ multiplet structure (horizontal in our diagrams). For the weight 2 fields for the $m = 1$ theory, this is the T_2 tilting module (or its Morita equivalent in the supersymmetric theories [20, 22]), which has the trio form

$$\begin{array}{ccc} 0 & & \\ & \searrow & \\ & 2 & \\ & \swarrow & \\ 0 & & \end{array} \quad (6.7)$$

in terms of the spin labels which are $j = 0, 2$ (for the $m = 1$ theory). This is in agreement with their expectations, since our node $j = 2$ is a multiplet, not a single field. What plays the role of supersymmetry in our case is $U_q(\mathfrak{sl}_2)$ or the $\mathcal{A}_{n+1|n}$ symmetry algebra [22], which is represented acting horizontally on Fig. 8. The dimension of the space of fields of conformal weight two in our supersymmetric models turns out to be:

$$\#\{\text{fields with } h = 2\} = D'_0 + D'_2 = (2n + 1)^4 - 3(2n + 1)^2 + 1 \quad (6.8)$$

(For the limit of the spin-1/2 chain, it is $1 + 5 = 6$.) The basic doublet t, T is thus completed by a large number (the preceding number minus 2, called D'_2) of other fields with $h = 2$, which themselves lie in the R_2 piece of another Virasoro indecomposable of structure

$$\mathcal{R}_3 : \quad \begin{array}{ccc} & R_3 & \\ \swarrow & & \searrow \\ R_2 & & R_5 \\ \searrow & & \swarrow \\ & R_3 & \end{array} . \quad (6.9)$$

The whole picture is thus considerably more involved than in Ref. [16, 17, 18].

A similar discussion applies to the dilute polymer case. The Virasoro tilting module containing the identity and the stress tensor is then

$$\mathcal{R}_1 : \quad \begin{array}{ccc} & R_1 & \\ \swarrow & & \searrow \\ R_0 & & R_2 \\ \searrow & & \swarrow \\ & R_1 & \end{array} \quad (6.10)$$

and the arrow from R_1 to R_2 indicates that in this case the OPE (6.1) must be extended to include a field of dimension 7, similar to τ ; it is a Virasoro descendant that is a null vector, but must not be set to zero. We have for the supersymmetric models

$$\#\{\text{fields with } h = 2\} = D'_0 + D'_1 = 4n^2. \quad (6.11)$$

For the $U_q(\mathfrak{sl}_2)$ theory, this number is $1 + 3 = 4$, so that it indeed turns out to be simply a quartet of fields, and as $q = i$ these can be viewed as a supersymmetry multiplet also, by putting $n = 1$.

It is interesting to observe that reducible but indecomposable modules of Virasoro of the forms we find are in general characterized by dimensionless parameters measuring the “mixing” of representations, and that the number b of Gurarie [16] appears simply as a particular case of what might be an infinite set of parameters. Clearly, a more detailed analysis of indecomposables at $c = 0$ along the lines of [30] would be useful here. But it is also important to understand how features of the lattice models determine the values of these parameters in our cases.

Gurarie and Ludwig [16, 17, 18] also advocated viewing the operator product algebra generated by t and T as a kind of chiral algebra. But we now see that we must introduce τ as a further field, and the fusion rules for this multiplet with itself produce fields in higher- j projective modules. That is, they generate by repeated ope’s not only the full set of integer-weight fields, which would then need to be included in the chiral algebra (and which are not all singlets under the symmetry algebra), but also for $m = 1$ the non-integer weight fields. In the oriented $m = 0$ and $m = 1$ theories, this space of fields corresponds to the full space of states! This argument ignored the role of the symmetry algebra in the actual fields, which after all carry representations of the symmetry algebra also. This might restrict the fields actually produced in iterated ope’s. However, it seems unfair to omit the fields related to t and T by symmetry from such a chiral algebra. The fields related by symmetry include some in the $\mathcal{R}_3$ Virasoro projective module, and fusion of these produces higher j ’s. Thus it seems inescapable that all integer j fields, including those with fractional weights, must be in such a chiral algebra. In cases such as the dilute unoriented polymers, there are also fields with $2j$ odd, and all of these together should probably be viewed as making up a single representation of this large chiral algebra. (Thus, the

structure of the modules over this larger algebra resembles that of the level-one $\mathfrak{su}(2)$ affine Lie algebra theory at $c = 1$, of which the $U_q(\mathfrak{sl}_2)$ theories are a deformation.) To us, it does not seem likely to be helpful to try to view this as a chiral algebra. Instead, the maximal chiral algebra (in the strict sense of containing only integer-weight fields) of these boundary theories seems to be the Virasoro algebra. On the other hand, the symmetry analysis seems powerful.

Finally, we turn to another issue: we point out that neither of the boundary CFTs with $c = 0$ possesses any fields with $h = 1$. The bulk theories in the supersymmetric formulation do possess such fields of weights $(1, 0)$ and $(0, 1)$ [20], and they transform in the adjoint representation of the algebras $\mathfrak{gl}(1 + n|n)$ and $\mathfrak{osp}(2n|2n)$, respectively in the two examples. The existence of these fields is required by Noether’s theorem. (Despite the conformal weights, these currents are not generally (anti-)holomorphic, implying that logarithms will appear in their correlators, and they do not generate affine Lie algebras [20]). In the boundary theory, on the other hand, while there are global symmetries, there is in general no such thing as a divergenceless boundary current, because charge can escape into the bulk, and Noether’s theorem only implies the existence of bulk currents. Thus in general there need not be any corresponding weight-1 fields in the boundary CFT.

On the other hand, a more fruitful point of view about such fields is that weight one fields on the boundary correspond to marginal perturbations of the boundary theory, while boundary fields of weight less than one are relevant perturbations on the boundary. In particular, exactly marginal perturbations of the boundary theory correspond to the possibility of a continuous change of conformally-invariant boundary conditions. For the $c = 0$ theories studied here, we learn that there are no marginal boundary perturbations, and the only relevant perturbations transform in representations of the symmetry algebra that do not contain any singlets. Thus such a perturbation would necessarily break the global symmetry.

On the other hand, for the $c = -2$ theories, there are a number of weight 1 fields, transforming in a T_2 tilting module for $U_q(\mathfrak{sl}_2)$, or a corresponding tilting module for $\mathcal{A}_{n|n}$. For symplectic fermions, there are 8 such fields. One of the fields in the $j = 2$ subquotient is the boundary term that corresponds to the bulk “topological” term in the sigma model, and corresponds to changing the Hall angle in the boundary condition discussed in Sec. 3. This gives a line of boundary conditions that preserve conformal invariance on the boundary. However, as we noted above, these theories can be transformed onto one another by a duality rotation, so this perturbation is “redundant”. The behavior under all the other perturbations is currently not known to us.

7 Conclusion

The analysis we have presented can be extended to the case of other values of q a root of unity, in relation with loop models. It can also be extended to other boundary conditions for open chains, which lead generically to standard modules with $h_{r,s}$, $r > 1$ using results from Refs. [48] and [49]; see also Ref. [23]. In all cases, the patterns are quite similar, in particular the reducible indecomposable projective Virasoro modules usually have the quartet form.

The real problem now is to extend the construction to the closed or periodic boundary condition case: we hope to report on this elsewhere.

While this paper and its companion were being completed, an interesting preprint appeared on the archive [50] which has some overlap with our analysis of the dense polymers case. In this reference, some of the fusion rules of the conformal field theory are conjectured based on a diagrammatic interpretation of fusion within the TL algebra. Our paper provides the correct algebraic framework to understand the observations of [50], together with the tools allowing immediate generalization to all roots of unity cases. We also would like to mention the recent work [51] where $U_q(\mathfrak{sl}_2)$ at roots of unity is used to build indecomposable representations of chiral W -algebras. As discussed in [20] the theories we are considering are not rational (though most likely quasi-rational) and we do not think the results of [51] apply to our case.

Acknowledgments

N.R. is grateful to the NSF for support under grant number DMR-02-42949. H.S. thanks V. Schomerus for useful discussions.

References

- [1] M. Flohr, Int. J. Mod. Phys. A **18**, 4497 (2003); M. Gaberdiel, Int. J. Mod. Phys. A **18**, 4593 (2003).
- [2] M. Flohr and A. Mueller-Lohmann, J. Stat. Mech. **0604**, P002 (2006).
- [3] I.A. Gruzberg, A.W.W. Ludwig, and N. Read, Phys. Rev. Lett. **82**, 4524 (1999).
- [4] S. Guruswamy, A. Leclair and A.W.W. Ludwig, Nucl. Phys. B **583**, 475 (2000).
- [5] M.J. Bhaseen, J.-S.Caux, I.I. Kogan and A.M. Tsvelik, Nucl. Phys. B **618**, 465 (2001).
- [6] G. Gotz, T. Quella and V. Schomerus, “The WZNW model on $PSU(1,1|2)$ ”, hep-th/0610070.
- [7] V.G. Knizhnik, Comm. Math. Phys. **112**, 567 (1987).
- [8] L. Rozansky and H. Saleur, Nucl. Phys. B **376**, 461 (1992).
- [9] V. Gurarie, Nucl. Phys. B **410**, 535 (1993).
- [10] M.R. Gaberdiel and H.G. Kausch, Phys. Lett. B **386**, 131 (1996); Nucl. Phys. B **538**, 631 (1999); H.G. Kausch, Nucl. Phys. B **583**, 513 (2000).
- [11] V. Schomerus and H. Saleur, Nucl. Phys. B **734**, 221 (2006).
- [12] M.A.I. Flohr, Int. J. Mod. Phys. A **11**, 4147 (1996); *ibid.* A **12**, 1943 (1997).
- [13] B. L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I.Yu Tipunin, Commun. Math. Phys. **265**, 47 (2006).
- [14] J. Fjelstad, J. Fuchs, S. Hwang, A.M. Semikhatov and I. Yu Tipunin, Nucl. Phys. B **633**, 379 (2002).
- [15] M. Flohr and A. Müller-Lohmann, hep-th/0510096.
- [16] V. Gurarie, Nucl. Phys. B **546**, 765 (1999).
- [17] V. Gurarie and A. W. W. Ludwig, J. Phys. A **35**, L377 (2002).
- [18] V. Gurarie and A. W. W. Ludwig, hep-th/0409105.
- [19] J. Cardy, “The stress tensor in quenched random systems”, in *Statistical Field Theories*, Proceedings of the NATO Advanced Research Workshop on Statistical Field Theories, June 2001, A. Cappelli and G. Mussardo, Eds, NATO Science series, Vol. II/73 (Kluwer Academic, Dordrecht, 2002).
- [20] N. Read and H. Saleur, Nucl. Phys. B **613**, 409 (2001).
- [21] G. Moore and N. Seiberg, Commun. Math. Phys. **123**, 177 (1989).
- [22] N. Read and H. Saleur, “Enlarged symmetry algebras of spin chains, loop models, and S -matrices”, cond-mat/0701xxx.
- [23] P. A. Pearce, J. Rasmussen and J.B. Zuber, “Logarithmic minimal models”, hep-th/0607232.
- [24] P.P. Martin, *Potts Models and Related Problems in Statistical Mechanics* (World Scientific, Singapore, 1991).

- [25] P. P. Martin, Int. J. Mod. Phys. A **7**, Supp. 1B, 645 (1992).
- [26] P. P. Martin and D. McAnally, Int. J. Mod. Phys. A **7**, Supp. 1B, 675 (1992).
- [27] V.F.R. Jones, Invent. Math. **72**, 1 (1983).
- [28] W.M. Koo and H. Saleur, Nucl. Phys. B **426**, 459 (1994).
- [29] G. Feverati and P. Pearce, Nucl. Phys. B **663**, 409 (2003).
- [30] F. Rohsiepe, “On Reducible but Indecomposable Representations of the Virasoro Algebra”, hep-th/9611160.
- [31] F.W. Anderson and K.R. Fuller, *Rings and Categories of Modules*, 2nd Ed., Graduate Texts in Mathematics 13 (Springer-Verlag, New York, NY, 1992).
- [32] For a review, see e.g. P. Di Francesco, P. Mathieu, and D. Senechal, *Conformal Field Theory* (Springer, New York, 1997).
- [33] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, Nucl. Phys. B **241**, 333 (1984).
- [34] H. Saleur and M. Bauer, Nucl. Phys. B **320**, 591 (1989).
- [35] J. Cardy, J. Stat. Phys. **125**, 1 (2006).
- [36] M.B. Green, J.H. Schwarz, and E. Witten, *Superstring Theory, Volume 2: Loop Amplitudes, Anomalies, and Phenomenology*, (Cambridge University, Cambridge, 1987), p. 204 ff.
- [37] B. Nienhuis, Phys. Rev. Lett. **49**, 1062 (1982).
- [38] P. di Francesco, H. Saleur, and J.B. Zuber, J. Stat. Phys. **49**, 57 (1987).
- [39] B. Nienhuis, J. Stat. Phys. **34**, 731 (1984).
- [40] V. Pasquier and H. Saleur, Nucl. Phys. B **330**, 523 (1990).
- [41] See Ref. [31], Lemma 29.4, p. 324.
- [42] S. Xiong, N. Read, and A.D. Stone, Phys. Rev. B **56**, 3982 (1997).
- [43] H.G. Kausch, “Curiosities at $c = -2$ ”, hep-th/9510149.
- [44] M. Gaberdiel and H. G. Kausch, “Indecomposable fusion products”, hep-th/9604026.
- [45] M.R. Gaberdiel, Int. J. Mod. Phys. A **9**, 4619 (1994).
- [46] P. di Francesco, H. Saleur and J.B. Zuber, Nucl. Phys. B **285**, 454 (1987).
- [47] U. Grimm and P. Pearce, J.Phys. A **26**, 7435 (1993).
- [48] P. Martin and H. Saleur, Lett. in Math. Phys. **30**, 189 (1994).
- [49] A. Nichols, “The Temperley-Lieb algebra and its generalizations in the Potts and XXZ models”, hep-th/0509069.
- [50] P. Pearce and J. Rasmussen, “Solvable critical dense polymers”, hep-th/0610273.
- [51] B. Feigin, A.M. Gainutdinov, A.M. Semikhatov and I. Yu Tipunin, “Kazhdan-Lusztig dual quantum group for logarithmic extensions of Virasoro minimal models”, hep-th/0606506.